arXiv:1210.4947v2 [math.DS] 20 Oct 2012

ROTOPULSATORS OF THE CURVED N-BODY PROBLEM

Florin Diacu[1,2] and **Shima Kordlou**[2]
[1]Pacific Institute for the Mathematical Sciences
and
[2]Department of Mathematics and Statistics
University of Victoria
P.O. Box 3060 STN CSC
Victoria, BC, Canada, V8W 3R4
diacu@uvic.ca and shimak@uvic.ca

October 23, 2012

Abstract. We consider the N-body problem in spaces of constant curvature and study its rotopulsators, i.e. solutions for which the configuration of the bodies rotates and, usually, changes size during the motion. Rotopulsators fall naturally into five groups: positive elliptic, positive elliptic-elliptic, negative elliptic, negative hyperbolic, and negative elliptic-hyperbolic, depending on the nature and number of their rotations and on whether they occur in spaces of positive or negative curvature. After obtaining existence criteria for each type of rotopulsator, we derive their conservation laws. We further deal with the existence and uniqueness of some classes of rotopulsators in the 3-body case and prove two general results about the qualitative behaviour of these orbits. An interesting finding is that of a class of rotopulsators that behave like relative equilibria, i.e. maintain constant mutual distances during the motion, but cannot be generated from any single element of the underlying subgroup $SO(2) \times SO(2)$ of the Lie group $SO(4)$.

1. Introduction

The goal of this paper is to study rotopulsators of the curved N-body problem, i.e. a type of solution that extends the concept of homographic orbit from the Euclidean space to spaces of constant curvature. In curved spaces, similar geometric figures are also congruent, so the word homographic is not suited for describing orbits that rotate and expand or contract during the motion. We will therefore introduce the concept of rotopulsating orbit, which coincides with the notion of homographic solution when the curvature of the space becomes zero. But before describing the results we prove in this paper, let us give the overall motivation of this research and provide a brief history of the N-body problem in spaces of constant curvature.

2. Motivation

The curved N-body problem offers an opportunity to look into the nature of the physical space. How do we measure the shortest distance between two points: along a straight line (like on a flat table), along an arc of a great circle (as between Paris and New York), or along a geodesic of some other manifold? Apparently, Carl Friedrich Gauss tried to answer this question by measuring the angles of triangles formed by three mountain peaks, to decide whether their sum was smaller or larger than π radians. But his experiments failed because no deviation from π could be detected outside the unavoidable measurement errors. It thus became clear that if the physical space is not flat, the deviation from zero curvature is very small. Some recent cosmological experiments involving the background radiation made physicists believe that space is Euclidean, although they have no proof so far, their results being at this point as inconclusive as those of Gauss. In fact, from the mathematical point of view, zero curvature is highly unlikely, if compared to a continuum of possibilities for positive or negative curvature.

The study of the curved N-body problem, however, may offer some new insight in this direction. We cannot travel to distant stars to measure angles of triangles, but we can observe celestial motions from Earth. If we prove the existence of orbits that characterize only one of the positive, zero, or negative constant curvature spaces, i.e. don't occur in any of the two others, and are stable too, then we can hope to decide the shape of the physical space by astronomical observations.

This new approach towards understanding the shape of the universe is likely to succeed at the local, but not the global, level because celestial motions at large scales are mainly radial: galaxies and clusters of galaxies merely move away from each other. Interesting celestial orbits occur only in solar systems. But the study of the curved N-body problem offers a new mathematical play field that can shed some light on the Euclidian case, by studying the bifurcations that occur when the curvature tends to zero, and may lead to a better understanding of several mathematical questions, including that of singularities in the motion of particle systems, [12], [4].

3. A brief history of the curved N-body problem

In the 1830s, János Bolyai and Nikolai Lobachevsky independently proposed a 2-body problem in the hyperbolic space $\mathbb{H}^3$, [2], [17]. They suggested the use of a force that is inversely proportional with the area of a sphere of radius r, where r is the distance between the bodies. Their line of thought followed that of Newton, who had viewed gravitation as inversely proportional to the area of an Euclidean sphere. But neither Bolyai nor Lobachevsky came up with an analytic expression for this new force. In 1870, Ernest Schering pointed out that, in $\mathbb{H}^3$, the area of a sphere of radius r is proportional to $\sinh^2 r$, so he defined a potential that involves

$\coth r$, [21]. Wilhelm Killing naturaly extended this idea to the sphere $\mathbb{S}^3$ with the help of a potential proportional to $\cot r$, [13].

In the first years of the 20th century, Heinrich Liebmann studied the Kepler problem (the motion of a body around a fixed centre) for the cotangent potential and recovered Kepler's laws in slightly modified form, [15]. Moreover, he found that a property earlier proved by Joseph Bertrand for the Newtonian potential, namely that all bounded orbits of the Kepler problem are closed, [1], was also true for the cotangent potential, [16]. These results, together with the fact that both the Newtonian and the cotangent potential of the Kepler problem are harmonic functions in the 3-dimensional case, established the cotangent potential among the problems worth researching in celestial mechanics. More recently, results in this direction were obtained by several Spanish and Russian mathematicians, [3], [14], [22]. It is interesting to note that, unlike in Euclidean space, the curved Kepler problem and the curved 2-body problem are not equivalent. While the former is integrable, the latter is not, [22], so its study appears to be complicated than that of the classical case.

Recently, we found a new setting that allowed the generalization of the 2-body case to any number $N \geq 2$ of bodies, [11], [6], [7], where we showed that the equations can be simultaneously written for positive and negative curvature. The idea was to use the hyperbolic sphere, i.e. Weierstrass's model of hyperbolic geometry, given by the upper sheet of the hyperboloid of two sheets, embedded in the Minkowski space. By suitable coordinate and time transformations, this study can be reduced to $\mathbb{S}^3$ and $\mathbb{H}^3$, or to $\mathbb{S}^2$ and $\mathbb{H}^2$ in the 2-dimensional case. For the latter, the equations in intrinsic coordinates were also obtained, [20], [10]. So far, we studied the singularities of the equations and of the solutions, the various classes of relative equilibria, and some rotopulsating orbits in the 2-dimensional case, [4], [5], [6], [7], [9], [10], [11], [12]. This paper provides a first investigation of the rotopulsating orbits in $\mathbb{S}^3$ and $\mathbb{H}^3$.

4. RESULTS

The remaining part of this paper is structured as follows. We first introduce in Section 5 the equations of motion in $\mathbb{S}^3$ and $\mathbb{H}^3$ and obtain their seven first integrals: one for the energy and six for the total angular momentum. In Section 6, we define the rotopulsating orbits (or rotopulsators) and classify them into five types, which we call positive elliptic, positive elliptic-elliptic, negative elliptic, negative hyperbolic, and negative elliptic-hyperbolic, depending on the manifold in which they occur and on the nature and number of their rotations.

The main results occur in Sections 7 through 13. Sections 7, 8, 10, 11, and 12 have the same structure, each providing a criterion for the existence of the type of rotopulsator it analyzes, giving the corresponding conservation laws, and proving existence and uniqueness results for classes of such orbits in the case $N = 3$. Thus

Section 7 deals with positive elliptic, Section 8 with positive elliptic-elliptic, Section 10 with negative elliptic, Section 11 with negative hyperbolic, and Section 12 with negative elliptic-hyperbolic rotopulsators. The examples provided in each of these sections are of Lagrangian type (i.e. the bodes lie at the vertices of rotating equilateral triangles) for the positive elliptic, positive elliptic-elliptic, and negative elliptic rotopulsators, and of Eulerian type (i.e. the bodies lie on the same rotating geodesic) for the negative hyperbolic and negative elliptic-hyperbolic rotopulsators.

Sections 10 and 13 offer a theorem each, whose object is to describe the qualitative behaviour of rotopulsators in $\mathbb{S}^3$ and $\mathbb{H}^3$, respectively. The first theorem shows that, for rotopulsators of $\mathbb{S}^3$, for any foliation of the sphere with Clifford tori, none of the bodies can stay confined to some Clifford torus, but each body passes through a continuum of such surfaces. The second theorem proves a similar result for $\mathbb{H}^3$, where the Clifford tori are replaced by hyperbolic cylinders.

An interesting finding is that of a class of Lagrangian orbits (see Subsection 8.3), which behave like relative equilibria, in the sense that they maintain constant mutual distances during the motion, but cannot be generated from any single element of the underlying subgroup $SO(2) \times SO(2)$ of the Lie group $SO(4)$. We call these rotopulsating orbits nonstandard relative equilibria.

5. EQUATIONS OF MOTION

Consider N point particles (bodies) of masses $m_1, \dots, m_N > 0$ moving in $\mathbb{S}^3$ (thought as embedded in the Euclidean space $\mathbb{R}^4$) or $\mathbb{H}^3$ (embedded in the Minkowski space $\mathbb{R}^{3,1}$), where

$$\mathbb{S}^3 = \{(w, x, y, z) \mid w^2 + x^2 + y^2 + z^2 = 1\},$$

$$\mathbb{H}^3 = \{(w, x, y, z) \mid w^2 + x^2 + y^2 - z^2 = -1, \ z > 0\}.$$

In previous work, we derived the equations of motion of the curved N-body problem using constrained Lagrangian dynamics and showed that by suitable coordinate and time-rescaling transformations the study of the problem can be reduced to $\mathbb{S}^3$ and $\mathbb{H}^3$, [6], [7].

The configuration of the system is described by the $4N$-dimensional vector

$$\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N),$$

where $\mathbf{q}_i = (w_i, x_i, y_i, z_i), i = 1, \dots, N$, denote the position vectors of the bodies. The equations of motion are given by the second-order system

$$\ddot{\mathbf{q}}_i = \sum_{j=1, j\neq i}^{N} \frac{m_j[\mathbf{q}_j - \sigma(\mathbf{q}_i \odot \mathbf{q}_j)\mathbf{q}_i]}{[\sigma - \sigma(\mathbf{q}_i \odot \mathbf{q}_j)^2]^{3/2}} - \sigma(\dot{\mathbf{q}}_i \odot \dot{\mathbf{q}}_i)\mathbf{q}_i, \quad i = 1, \dots, N, \tag{1}$$

with constraints

$$\mathbf{q}_i \odot \mathbf{q}_\mathbf{i} = \sigma, \quad \mathbf{q}_i \odot \dot{\mathbf{q}}_i = 0, \quad i = 1, \dots, N. \tag{2}$$

Here $\odot$ is the standard inner product $\cdot$ of signature $(+,+,+,+)$ in $\mathbb{S}^3$ or the Lorentz inner product $\boxdot$ of signature $(+,+,+,-)$ in $\mathbb{H}^3$, and

$$\sigma = \begin{cases} +1, & \text{in } \mathbb{S}^3, \\ -1, & \text{in } \mathbb{H}^3, \end{cases}$$

denotes the signum function. Equations (1) can be viewed as a $6N$-dimensional first-order system of ordinary differential equations. The force acting on each body has a gravitational component (the above sum) and a term (involving the velocities) that corresponds to the constraints.

As a consequence of Noether's theorem, system (1) has the scalar integral of energy,

$$T(\mathbf{q}, \dot{\mathbf{q}}) - U(\mathbf{q}) = h,$$

where

$$U(\mathbf{q}) = \sum_{1 \le i < j \le N} \frac{\sigma m_i m_j \mathbf{q}_i \odot \mathbf{q}_j}{[\sigma - \sigma(\mathbf{q}_i \odot q_j)^2]^{3/2}}$$

is the force function ($-U$ representing the potential), which stems from the cotangent of the distance,

$$T(\mathbf{q}, \dot{\mathbf{q}}) = \frac{1}{2} \sum_{i=1}^{N} m_i (\dot{\mathbf{q}}_i \odot \dot{\mathbf{q}}_i)(\sigma \mathbf{q}_i \odot \mathbf{q}_i)$$

is the kinetic energy, with h representing an integration constant. System (1) also has the 6-dimensional integral of the total angular momentum,

$$\sum_{i=1}^{N} m_i \mathbf{q}_i \wedge \dot{\mathbf{q}}_i = \mathbf{c},$$

where $\wedge$ is the wedge product and $\mathbf{c} = (c_{wx}, c_{wy}, c_{wz}, c_{xy}, c_{xz}, c_{yz})$ denotes an integration vector, each component measuring the rotation of the system about the origin of the frame relative to the plane corresponding to the bottom indices.

For positive curvature, i.e. in $\mathbb{S}^3$, we can explicitly write the equations of motion as

$$\ddot{\mathbf{q}}_i = \sum_{\substack{j=1 \\ j \ne i}}^{N} \frac{m_j [\mathbf{q}_j - (\mathbf{q}_i \cdot \mathbf{q}_j)\mathbf{q}_i]}{[1 - (\mathbf{q}_i \cdot \mathbf{q}_j)^2]^{3/2}} - (\dot{\mathbf{q}}_i \cdot \dot{\mathbf{q}}_i)\mathbf{q}_i, \;\; \mathbf{q}_i \cdot \mathbf{q_i} = 1, \quad i = 1, 2, \dots, N, \tag{3}$$

while for negative curvature, i.e. in $\mathbb{H}^3$, they take the form

$$\ddot{\mathbf{q}}_i = \sum_{\substack{j=1 \\ j \ne i}}^{N} \frac{m_j [\mathbf{q}_j + (\mathbf{q}_i \boxdot \mathbf{q}_j)\mathbf{q}_i]}{[(\mathbf{q}_i \boxdot \mathbf{q}_j)^2 - 1]^{3/2}} + (\dot{\mathbf{q}}_i \boxdot \dot{\mathbf{q}}_i)\mathbf{q}_i, \;\; \mathbf{q}_i \boxdot \mathbf{q_i} = -1, \quad i = 1, 2, \dots, N. \tag{4}$$

Systems (3) and (4) are the forms of the equations of motion we will mainly refer to in the remainder of this paper.

6. CLASSIFICATION

In this section we define several types of rotopulsating orbits of the curved N-body problem and explain the choices we make to classify them. The rotopulsating orbit extends the Euclidean homographic solution to spaces of nonzero constant curvature, which actually reduce to the unit 3-sphere $\mathbb{S}^3$, in the positive case, and to the unit hyperbolic sphere, $\mathbb{H}^3$, for negative curvature. In two previous papers, we introduced this concept in the 2-dimensional case and kept using the name "homographic" for it, [5], [9]. Our idea was that the configurations we studied (mostly polygons) remained homographic if viewed in the ambient Euclidean space. But it seems more natural to regard configurations in intrinsic terms, the more so when we move from two to three dimensions.

In $\mathbb{S}^2, \mathbb{S}^3, \mathbb{H}^2$, and $\mathbb{H}^3$, however, the concept of similarity, which corresponds to the adjective homographic, makes little sense, since, for instance, the only similar triangles are the congruent ones. Consequently, to extend the concept of homographic orbit to spaces of constant curvature, we need, on one hand, to capture somehow the dilation/contraction aspect as well as the rotation component of the solutions, and, on the other hand, to recover the original definition when the curvature tends to zero. We therefore introduce here a new adjective, *rotopulsating*, which suggests both these features of the orbit without implying similarity of the configuration. For rotopulsating orbits will be also use the noun *rotopulsator*.

The definitions we provide below follow naturally from the concept of relative equilibrium of the curved N-body problem, defined in [6], [7]. We introduced there various kinds of relative equilibria in terms of the isometric rotation groups of $\mathbb{S}^3$ and $\mathbb{H}^3$. The rotopulsating orbits we will further define differ from the relative equilibria through nonuniform rotations and, in general, through nonconstant mutual distances, as we will further see.

Definition 1 (**Positive elliptic rotopulsators).** *A solution of system* (3) *in* $\mathbb{S}^3$ *is called a positive elliptic rotopulsator if it is of the form*

$$\begin{gathered}\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, \dots, N, \\ w_i = r_i(t)\cos[\alpha(t) + a_i], \quad x_i = r_i(t)\sin[\alpha(t) + a_i], \quad y_i = y_i(t), \quad z_i = z_i(t),\end{gathered} \tag{5}$$

where $a_i,\ i = 1, 2, \dots, N$, *are constants,* α *is a nonconstant function,* r_i, y_i, *and* z_i *satisfy the conditions*

$$0 \le r_i \le 1, \quad -1 \le y_i, z_i \le 1, \quad \text{and} \quad r_i^2 + y_i^2 + z_i^2 = 1, \quad i = 1, 2, \dots, N,$$

and $c_{yz} = 0$. *If* r *is constant and* $\alpha(t) = \bar{\alpha}t$, *with* $\bar{\alpha}$ *a nonzero constant, then the solution is called a positive elliptic relative equilibrium.*

Remark 1. The conditions that α is nonconstant and $c_{yz} = 0$ are imposed to ensure that the system has an elliptic rotation relative to the wx-plane, but no elliptic rotation relative to the yz-plane. Rotations relative to other base planes may occur.

Definition 2 (**Positive elliptic-elliptic rotopulsators**). *A solution of system* (3) *in* $\mathbb{S}^3$ *is called a positive elliptic-elliptic rotopulsator if it is of the form*

$$\begin{gathered}\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N),\ \mathbf{q}_i = (w_i, x_i, y_i, z_i),\ i = 1, 2, \dots, N,\\ w_i = r_i(t)\cos[\alpha(t) + a_i],\ x_i = r_i(t)\sin[\alpha(t) + a_i],\\ y_i = \rho_i(t)\cos[\beta(t) + b_i],\ z_i = \rho_i(t)\sin[\beta(t) + b_i],\end{gathered} \tag{6}$$

where $a_i, b_i,\ i = 1, 2, \dots, N,$ *are constants,* α *and* β *are nonconstant functions, and* r_i *and* ρ_i *satisfy the conditions*

$$0 \le r_i, \rho_i \le 1 \ \text{ and } \ r_i^2 + \rho_i^2 = 1,\ i = 1, 2, \dots, N.$$

If r *and* ρ *are constant and* $\alpha(t) = \bar{\alpha}t, \beta(t) = \bar{\beta}t$, *with* $\bar{\alpha}, \bar{\beta}$ *nonzero constants, then the solution is called a positive elliptic-elliptic relative equilibrium.*

Remark 2. The condition that α and β are nonconstant is imposed to ensure that the system has two elliptic rotations, one relative to the wx-plane and the other relative to the yz-plane. Rotations relative to the other base planes may occur.

Definition 3 (**Negative elliptic rotopulsators**). *A solution of system* (4) *in* $\mathbb{H}^3$ *is called a negative elliptic rotopulsator if it is of the form*

$$\begin{gathered}\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, \dots, N,\\ w_i = r_i(t)\cos[\alpha(t) + a_i], \quad x_i = r_i(t)\sin[\alpha(t) + a_i], \quad y_i = y_i(t), \quad z_i = z_i(t),\end{gathered} \tag{7}$$

where $a_i,\ i = 1, 2, \dots, N,$ *are constants,* α *is a nonconstant function,* $r_i, y_i,$ *and* z_i *satisfy the conditions*

$$z_i \ge 1 \quad \text{and} \quad r_i^2 + y_i^2 - z_i^2 = -1, \quad i = 1, 2, \dots, N,$$

and $c_{yz} = 0$. *If* r *is constant and* $\alpha(t) = \bar{\alpha}t$, *with* $\bar{\alpha}$ *a nonzero constant, then the solution is called a negative elliptic relative equilibrium.*

Remark 3. The conditions that α is nonconstant and $c_{yz} = 0$ are imposed to ensure that the system has an elliptic rotation relative to the wx-plane, but no hyperbolic rotation relative to the yz-plane. Rotations relative to other base planes may occur.

Definition 4 (**Negative hyperbolic rotopulsators**). *A solution of system* (4) *in* $\mathbb{H}^3$ *is called a negative hyperbolic rotopulsator if it is of the form*

$$\begin{gathered}\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, \dots, N,\\ w_i = w_i(t),\ x_i = x_i(t),\ y_i = \eta_i(t)\sinh[\beta(t) + b_i],\ z_i = \eta_i(t)\cosh[\beta(t) + b_i],\end{gathered} \tag{8}$$

where $b_i,\ i=1,2,\dots,N,$ *are constants,* β *is a nonconstant function,* $w_i, x_i, z_i,$ *and* η_i *satisfy the conditions*

$$z_i \ge 1 \quad \text{and} \quad w_i^2 + x_i^2 - \eta_i^2 = -1, \quad i=1,2,\dots,N,$$

and $c_{wx}=0$*. If* η *is constant and* $\beta(t)=\bar\beta t$*, with* $\bar\beta$ *a nonzero constant, then the solution is called a negative hyperbolic relative equilibrium.*

Remark 4. The conditions that β is nonconstant and $c_{wx}=0$ are imposed to ensure that the system has a hyperbolic rotation relative to the yz-plane, but no elliptic rotation relative to the wx-plane. Rotations relative to other base planes may occur.

Definition 5 (**Negative elliptic-hyperbolic rotopulsators**)**.** *A solution of system* (4) *in* $\mathbb{H}^3$ *is called a negative elliptic-hyperbolic rotopulsator if it is of the form*

$$\begin{aligned} &\mathbf{q}=(\mathbf{q}_1,\mathbf{q}_2,\dots,\mathbf{q}_N), \quad \mathbf{q}_i=(w_i,x_i,y_i,z_i), \quad i=1,2,\dots,N,\\ &w_i=r_i(t)\cos[\alpha(t)+a_i], \quad x_i=r_i(t)\sin[\alpha(t)+a_i],\\ &y_i=\eta_i(t)\sinh[\beta(t)+b_i], \ z_i=\eta_i(t)\cosh[\beta(t)+b_i], \end{aligned} \tag{9}$$

where $a_i, b_i,\ i=1,2,\dots,N,$ *are constants,* α *and* β *are nonconstant functions, whereas* $r_i, \eta_i,$ *and* z_i *satisfy the conditions*

$$z_i \ge 1 \quad \text{and} \quad r_i^2 - \eta_i^2 = -1, \quad i=1,2,\dots,N.$$

When r *and* η *are constant and* $\alpha(t)=\bar\alpha t, \beta(t)=\bar\beta t$*, with* $\bar\alpha, \bar\beta$ *nonzero constants, then the solution is called a negative elliptic-hyperbolic relative equilibrium.*

Remark 5. The conditions for α and β are imposed to ensure that the system has an elliptic rotation relative to the wx-plane and a hyperbolic rotation relative to the yz-plane. Rotations relative to other base planes may occur.

Remark 6. There are alternative ways to define the above classes of rotopulsating solutions of the curved N-body problem (including those of considering positive elliptic orbits as particular positive elliptic-elliptic orbits, and negative elliptic and negative hyperbolic orbits as particular negative elliptic-hyperbolic orbits), but our choice is more convenient for computations and will help us emphasize, unambiguously, certain properties of all these solutions.

Before starting to study these orbits, let us remark that, in spaces of nonzero constant curvature, the main difference between relative equilibria and rotopulsators is the same we find in Euclidean space between relative equilibria and homographic orbits that are not relative equilibria. Namely, in the former class of orbits the mutual distances are constant in time, whereas in the latter class, the geometric configuration of the particle system changes in size, in general. This is the reason why we used in [6] and [7] the isometric rotation groups of $\mathbb{S}^3$ and $\mathbb{H}^3$

to define the relative equilibria, a move that led us naturally to five classes of such orbits. Allowing then the size of these configurations to vary, and implicitly making the rotation of the system nonuniform, we were led to the above definitions for rotopulsators.

In 3-dimensional spaces of constant curvature, however, a new possibility occurs, which does not take place in $\mathbb{R}^2, \mathbb{R}^3, \mathbb{S}^2$, and $\mathbb{H}^2$. Namely, there could exist rotopulsating orbits, as defined above, which maintain constant mutual distances. In other words, such motions (if they exist, and we will show that they do) are relative equilibria, though they don't belong to the class of orbits obtained from isometric rotations, i.e. do not obey the definitions of relative equilibria given above, or in [6] and [7]. Therefore the following new definition becomes necessary.

Definition 6. *Any of the rotopulsators defined in* (5)*,* (6)*,* (7)*,* (8)*, or* (9) *are called nonstandard relative equilibria if they are not relative equilibria in the sense of Definitions* 1*,* 2*,* 3*,* 4*, or* 5*, although the mutual distances between the bodies remain constant during the motion.*

As we will see later, these orbits occur due to the fact that they cannot be generated from a single element of the Lie group $SO(2) \times SO(2)$, which occurs naturally when describing such solutions. Nevertheless, they may occur from an element generated by $SO(4)$, as the more general theory of relative equilibria developed in the framework of geometric mechanics suggests, [19].

7. POSITIVE ELLIPTIC ROTOPULSATORS

In this section we analyze the solutions given in Definition 1. We first introduce a criterion for finding them, then obtain the conservation laws, and finally prove the existence of a particular class of orbits, namely the positive elliptic Lagrangian rotopulsators of the 3-body problem in $\mathbb{S}^3$.

7.1. Criterion for positive elliptic rotopulsators.

The following result provides necessary and sufficient conditions for the existence of positive elliptic rotopulsators in $\mathbb{S}^3$.

Criterion 1. *A solution candidate of the form* (5) *is a positive elliptic rotopulsator for system* (3) *if and only if* $c_{yz} = 0$*,*

$$\dot{\alpha} = \frac{c}{\sum_{j=1}^N m_j(1 - y_j^2 - z_j^2)}, \tag{10}$$

where c is a nonzero constant, and the variables $y_i, z_i,\ i = 1, 2, \dots, N,$ satisfy the system of $2N$ second-order differential equations

$$\begin{cases} \ddot{y}_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(y_j - q_{ij}y_i)}{(1-q_{ij}^2)^{3/2}} - G_i y_i \\ \ddot{z}_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(z_j - q_{ij}z_i)}{(1-q_{ij}^2)^{3/2}} - G_i z_i, \end{cases} \tag{11}$$

where

$$G_i := \frac{\dot y_i^2 + \dot z_i^2 - (y_i\dot z_i - z_i\dot y_i)^2}{1 - y_i^2 - z_i^2} + \frac{c^2(1 - y_i^2 - z_i^2)}{\left[\sum_{j=1}^N m_j(1 - y_j^2 - z_j^2)\right]^2}, \tag{12}$$

$i = 1, 2, \dots, N$, *and, for any* $i, j \in \{1, 2, \dots, N\}$, q_{ij} *is given by*

$$q_{ij} := \mathbf{q}_i \cdot \mathbf{q_j} = (1 - y_i^2 - z_i^2)^{\frac12}(1 - y_j^2 - z_j^2)^{\frac12}\cos(a_i - a_j) + y_iy_j + z_iz_j.$$

Proof. Consider a solution candidate of the form (5). A straightforward computation shows that, for any $i, j \in \{1, 2, \dots, N\}$, q_{ij} is of the form given in the above statement. Moreover, for any $i = 1, 2, \dots, N$, we find that

$$\dot{\mathbf{q}}_i \cdot \dot{\mathbf{q}}_i = \frac{\dot y_i^2 + \dot z_i^2 - (y_i\dot z_i - z_i\dot y_i)^2 + (1 - y_i^2 - z_i^2)^2\dot\alpha^2}{1 - y_i^2 - z_i^2}.$$

For all $i = 1, 2, \dots, N$, each r_i can be expressed in terms of y_i and z_i to obtain

$$r_i = (1 - y_i^2 - z_i^2)^{\frac12}, \quad \dot r_i = -\frac{y_i\dot y_i + z_i\dot z_i}{(1 - y_i^2 - z_i^2)^{\frac12}},$$

$$\ddot r_i = \frac{(y_i\dot z_i - z_i\dot y_i)^2 - \dot y_i^2 - \dot z_i^2 - (1 - y_i^2 - z_i^2)(y_i\ddot y_i + z_i\ddot z_i)}{(1 - y_i^2 - z_i^2)^{\frac32}}.$$

Substituting a candidate solution of the form (5) into system (3) and employing the above formulas, we obtain for the equations corresponding to $\ddot y_i$ and $\ddot z_i$ that

$$\ddot y_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(y_j - q_{ij}y_i)}{(1 - q_{ij}^2)^{\frac32}} - \frac{[\dot y_i^2 + \dot z_i^2 - (y_i\dot z_i - z_i\dot y_i)^2]y_i}{1 - y_i^2 - z_i^2} - (1 - y_i^2 - z_i^2)y_i\dot\alpha^2, \tag{13}$$

$$\ddot z_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(z_j - q_{ij}z_i)}{(1 - q_{ij}^2)^{\frac32}} - \frac{[\dot y_i^2 + \dot z_i^2 - (y_i\dot z_i - z_i\dot y_i)^2]z_i}{1 - y_i^2 - z_i^2} - (1 - y_i^2 - z_i^2)z_i\dot\alpha^2. \tag{14}$$

For the equations corresponding to $\ddot w_i$ and $\ddot x_i$, after some long computations that also use (13) and (14), we are led either to identities or to the equations

$$r_i\ddot\alpha + 2\dot r_i\dot\alpha = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_jr_j\sin(a_j - a_i)}{(1 - q_{ij}^2)^{\frac32}}, \quad i = 1, 2, \dots, N. \tag{15}$$

For every $i = 1, 2, \dots, N$, we further multiply the ith equation in (15) by m_ir_i, add the resulting N equations, and notice that

$$\sum_{i=1}^N \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_im_jr_ir_j\sin(a_j - a_i)}{(1 - q_{ij}^2)^{\frac32}} = 0.$$

Thus we obtain the equation

$$\ddot\alpha\sum_{i=1}^N m_ir_i^2+2\dot\alpha\sum_{i=1}^N m_ir_i\dot r_i=0,$$

which has the solution

$$\dot\alpha=\frac{c}{\sum_{i=1}^N m_ir_i^2}=\frac{c}{\sum_{i=1}^N m_i(1-y_i^2-z_i^2)},$$

where c is an integration constant. Consequently, equations (13) and (14) become

(16)
$$\ddot y_i=\sum_{\substack{j=1\\ j\neq i}}^N\frac{m_j(y_j-q_{ij}y_i)}{(1-q_{ij}^2)^{\frac32}}-\frac{[\dot y_i^2+\dot z_i^2-(y_i\dot z_i-z_i\dot y_i)^2]y_i}{1-y_i^2-z_i^2}-\frac{c^2(1-y_i^2-z_i^2)y_i}{\left[\sum_{j=1}^N m_j(1-y_j^2-z_j^2)\right]^2},$$

(17)
$$\ddot z_i=\sum_{\substack{j=1\\ j\neq i}}^N\frac{m_j(z_j-q_{ij}z_i)}{(1-q_{ij}^2)^{\frac32}}-\frac{[\dot y_i^2+\dot z_i^2-(y_i\dot z_i-z_i\dot y_i)^2]z_i}{1-y_i^2-z_i^2}-\frac{c^2(1-y_i^2-z_i^2)z_i}{\left[\sum_{j=1}^N m_j(1-y_j^2-z_j^2)\right]^2},$$

$i=1,2,\dots,N$. Since equations (16) and (17) lead to system (11), and the condition $c_{yz}=0$ occurs from Definition 1, the criterion is proved. □

7.2. **Conservation laws for positive elliptic rotopulsating orbits.** In addition to Criterion 1, we would also like to obtain the conservation laws specific to positive elliptic rotopulsators. They follow by straightforward computations using the above proof, the integral of energy, and the six integrals of the total angular momentum.

Proposition 1. *If system* (3) *has a solution of the form* (5), *then the following expressions are constant:*

— *energy,*

(18)
$$h=\sum_{i=1}^N\frac{m_i[\dot y_i^2+\dot z_i^2-(y_i\dot z_i-z_i\dot y_i)^2]}{2(1-y_i^2-z_i^2)}+\frac{c^2}{2\sum_{i=1}^N m_i(1-y_i^2-z_i^2)}-\sum_{1\le i<j\le N}\frac{m_im_jq_{ij}}{(1-q_{ij}^2)^{\frac12}};$$

— *total angular momentum relative to the wx-plane,*

(19)
$$c_{wx}=c,$$

where $c\neq0$ *is the constant in the expression* (15) *of* $\dot\alpha$*;*

— *total angular momentum relative to the wy-plane:*

$$\begin{aligned} c_{wy} &= \sum_{i=1}^N m_i\left[(1-y_i^2-z_i^2)^{\frac12}\dot y_i+\frac{(y_i\dot y_i+z_i\dot z_i)y_i}{(1-y_i^2-z_i^2)^{\frac12}}\right]\cos(\alpha+a_i)\\ &+\frac{c}{\sum_{i=1}^N m_i(1-y_i^2-z_i^2)}\sum_{i=1}^N m_i(1-y_i^2-z_i^2)^{\frac12}y_i\sin(\alpha+a_i); \end{aligned}\tag{20}$$

— *total angular momentum relative to the wz-plane,*

$$\begin{aligned} c_{wz} &= \sum_{i=1}^N m_i\left[(1-y_i^2-z_i^2)^{\frac12}\dot z_i+\frac{(y_i\dot y_i+z_i\dot z_i)z_i}{(1-y_i^2-z_i^2)^{\frac12}}\right]\cos(\alpha+a_i)\\ &+\frac{c}{\sum_{j=1}^N m_j(1-y_j^2-z_j^2)}\sum_{i=1}^N m_i(1-y_i^2-z_i^2)^{\frac12}z_i\sin(\alpha+a_i); \end{aligned}\tag{21}$$

— *total angular momentum relative to the xy-plane,*

$$\begin{aligned} c_{xy} &= \sum_{i=1}^N m_i\left[(1-y_i^2-z_i^2)^{\frac12}\dot y_i+\frac{(y_i\dot y_i+z_i\dot z_i)y_i}{(1-y_i^2-z_i^2)^{\frac12}}\right]\sin(\alpha+a_i)\\ &-\frac{c}{\sum_{j=1}^N m_j(1-y_j^2-z_j^2)}\sum_{i=1}^N m_i(1-y_i^2-z_i^2)^{\frac12}y_i\cos(\alpha+a_i); \end{aligned}\tag{22}$$

— *total angular momentum relative to the xz-plane:*

$$\begin{aligned} c_{xz} &= \sum_{i=1}^N m_i\left[(1-y_i^2-z_i^2)^{\frac12}\dot z_i+\frac{(y_i\dot y_i+z_i\dot z_i)z_i}{(1-y_i^2-z_i^2)^{\frac12}}\right]\sin(\alpha+a_i)\\ &-\frac{c}{\sum_{j=1}^N m_j(1-y_j^2-z_j^2)}\sum_{i=1}^N m_i(1-y_i^2-z_i^2)^{\frac12}z_i\cos(\alpha+a_i); \end{aligned}\tag{23}$$

— *total angular momentum relative to the yz-plane:*

$$c_{yz}=0.\tag{24}$$

The above result could be also used to prove the nonexistence of some candidates for positive elliptic rotopulsators, by showing that at least one of the above conservation laws is violated.

7.3. **Positive elliptic Lagrangian rotopulsators.** We further provide a class of specific examples of positive elliptic rotopulsators of the curved 3-body problem, namely Lagrangian orbits, i.e. bodies that lie at the vertices of a rotating equilateral triangle in $\mathbb{S}^3$ that changes size, which means that it fails to be similar to itself but has congruent sides at every instant in time. As we will see, these systems rotate relative to the wx-plane, but have no rotations with respect to the other base planes.

Consider three equal masses, $m_1 = m_2 = m_3 =: m$, and a candidate solution of the form

$$\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, 3, \tag{25}$$

$$w_1 = r(t)\cos\alpha(t),\ x_1(t) = r(t)\sin\alpha(t),\ y_1 = y(t),\ z_1(t) = z(t),$$

$$w_2 = r(t)\cos[\alpha(t) + 2\pi/3],\ x_2(t) = r(t)\sin[\alpha(t) + 2\pi/3], y_2 = y(t),\ z_2(t) = z(t),$$

$$w_3 = r(t)\cos[\alpha(t) + 4\pi/3],\ x_3(t) = r(t)\sin[\alpha(t) + 4\pi/3], y_3 = y(t),\ z_3(t) = z(t).$$

With the help of Criterion 1, we will further show that, in general, these are indeed solutions of system (3).

Proposition 2. *Consider the curved* 3*-body problem in* $\mathbb{S}^3$ *given by system* (3)*. Then, except for a negligible set of solutions formed by positive elliptic Lagrangian relative equilibria, every candidate solution of the form* (25) *is a positive elliptic Lagrangian rotopulsator, which rotates only relative to the* wx*-plane.*

Proof. Let us consider a candidate solution of the form (25). Then, using Criterion 1, straightforward computations show that

$$q_{12} = q_{21} = q_{13} = q_{31} = q_{23} = q_{32} = \frac{3y^2 + 3z^2 - 1}{2}, \quad \dot\alpha = \frac{c}{3m(1 - y^2 - z^2)},$$

and that the variables y and z must satisfy the equations

$$\begin{cases} \ddot y = F(y, z, \dot y, \dot z)y \\ \ddot z = F(y, z, \dot y, \dot z)z, \end{cases} \tag{26}$$

where

$$F(y, z, \dot y, \dot z) = \frac{8m}{\sqrt{3}(1 - y^2 - z^2)^{\frac12}(1 + 3y^2 + 3z^2)^{\frac32}} - \frac{c^2}{9m^2(1 - y^2 - z^2)}$$
$$- \frac{\dot y^2 + \dot z^2 - (y\dot z - \dot y z)^2}{1 - y^2 - z^2}.$$

From (26), we can conclude that $\ddot y z = y\ddot z$, which implies that

$$y\dot z - z\dot y = k \text{ (constant)}.$$

But, since from (24) we have that $3m(y\dot z - z\dot y) = c_{yz}$, it follows that $k = c_{yz}/3m$. As $c_{yz} = 0$, we must take $k = 0$, so $y\dot z - z\dot y = 0$, and therefore $\frac{d}{dt}\frac{y}{z} = 0$ if z does not take zero values, so $y(t) = \gamma z(t)$, where γ is a constant. Moreover, since

$$\sin\alpha + \sin(\alpha + 2\pi/3) + \sin(\alpha + 4\pi/3) = \cos\alpha + \cos(\alpha + 2\pi/3) + \cos(\alpha + 4\pi/3) = 0,$$

it follows from (20), (21), (22), and (23) that $c_{wy} = c_{wz} = c_{xy} = c_{xz} = 0$, so the triangle has no rotation relative to the planes wy, wz, xy, xz, and yz.

Notice that the energy relation (18) takes the form

$$\frac{3m[\dot y^2+\dot z^2-(y\dot z-\dot y z)^2]}{2(1-y^2-z^2)}+\frac{c^2}{6m(1-y^2-z^2)}-\frac{\sqrt{3}m^2(3y^2+3z^2-1)}{(1-y^2-z^2)^{\frac12}(1+3y^2+3z^2)^{\frac12}}=h,$$

which implies that F can be written as

$$F(y,z)=\frac{2m[5-9(y^2+z^2)^2]}{\sqrt{3}(1-y^2-z^2)^{\frac12}(1+3y^2+3z^2)^{\frac32}}-\frac{2h}{3m}.$$

If we denote $\delta=\gamma^2+1\ge 1$, system (26) reduces to the family of first order systems

$$\begin{cases}\dot z = v\\ \dot v=\left[\frac{2m(5-9\delta^2z^4)}{\sqrt{3}(1-\delta z^2)^{\frac12}(1+3\delta z^2)^{\frac32}}-\frac{2h}{3m}\right]z.\end{cases}\tag{27}$$

Standard results of the theory of ordinary differential equations can now be applied to system (27) to prove the existence and uniqueness of analytic positive elliptic Lagrangian rotopulsators, for admissible initial conditions. To show that only a negligible set of such orbits is given by relative equilibria, notice that finding the fixed points in the family of equations given by system (27), which depends on parameters, reduces to obtaining the zeroes of a polynomial equation. This remark completes the proof. □

7.4. **The nature of the solutions.** Although we proved the existence of positive elliptic rotopulsators, it would be still interesting to learn more about their nature. For this, we will first find the fixed points for the vector field of system (27) different from the obvious one, $(z,v)=(0,0)$, which is common to all equations in the family. This problem reduces to finding the zeroes of the family of polynomials

$$P(z)=27(9m^4+h^2)\delta^4z^8-18(15m^4+h^2)\delta^2z^4-8h^2\delta z^2+75m^4-h^2.$$

According to Descartes's rule of signs, we have to distinguish between two cases:

(i) $|h|<5\sqrt{3}m^2$, when, for every fixed values of the parameters, P has either two positive or no positive root at all;

(ii) $|h|\ge 5\sqrt{3}m^2$, when, for every fixed values of the parameters, P has exactly one positive root.

Case (ii) always leads to one fixed point since the unique positive root, $z_0 := z_0(m,h,\epsilon)$, is always such that $|z_0|<1$. To show this fact, let $\zeta=\delta z^2$. Then we can associate to P the family of polynomials

$$\bar P(\zeta)=27(9m^4+h^2)\zeta^4-18(15m^4+h^2)\zeta^2-8h^2\zeta+75m^4-h^2.$$

Since $\delta\ge 1$, to prove that $|z_0|<1$ it is enough show that $|\zeta_0|:=|\delta z_0|<1$. For this, notice that $\bar P(1)=48m^4>0$. As $\bar P(0)<0$ in case (ii), it follows that

$0 < \zeta_0 < 1$. Then the corresponding eigenvalues $\lambda_{1,2}$ are given by the equation

$$\lambda + \frac{2h}{3m} - W(z_0, m, \epsilon) = 0,$$

where $W(z_0, m, \epsilon)$ is a finite number for every fixed value of the parameters. Independently of the values of $W(z_0, m, \epsilon)$, the eigenvalues show that z varies for every orbit that is not a fixed point. Similar conclusions can be drawn in case (i).

Numerical experiments suggest that all the other orbits of system (27) are periodic, except for two homoclinic orbits. Since the periods of α and z don't usually match, the periodic orbits generate quasiperiodic positive elliptic Lagrangian orbits, except for a negligible class, given by periodic positive elliptic Lagrangian orbits (see Fig. 1).



FIGURE 1. The flow of system (27) for $m = 1, c_{yz} = 0, \delta = 1, h = 5$.

Remark 7. When y or z is a nonzero constant, the motion takes place on a 2-dimensional nongreat sphere. When $y \equiv 0$ or $z \equiv 0$, the motion takes place on a 2-dimensional great sphere, i.e. on $\mathbb{S}^2$, a type of Lagrangian orbit for which we gave a complete classification in [9].

8. POSITIVE ELLIPTIC-ELLIPTIC ROTOPULSATORS

In this section we analyze the solutions given in Definition 2. We first prove a criterion for finding such solutions, then obtain the conservation laws, and finally discuss a particular class of examples, namely the positive elliptic-elliptic Lagrangian rotopulsators of the 3-body problem in $\mathbb{S}^3$.

8.1. Criterion for positive elliptic-elliptic rotopulsators.

The following result provides necessary and sufficient conditions for the existence of positive elliptic-elliptic rotopulsators in $\mathbb{S}^3$.

Criterion 2. *A solution candidate of the form* (6) *is a positive elliptic-elliptic rotopulsator for system* (3) *if and only if*

$$\dot\alpha = \frac{c_1}{\sum_{i=1}^N m_i r_i^2}, \quad \dot\beta = \frac{c_2}{M - \sum_{i=1}^N m_i r_i^2}, \tag{28}$$

with $c_1, c_2, M = \sum_{i=1}^N m_i$ *nonzero constants, and the variables* $r_1, r_2, \dots, r_N$ *satisfy the* N *second-order differential equations*

$$\begin{aligned}\ddot r_i &= r_i(1-r_i^2)\left[\frac{c_1^2}{(\sum_{i=1}^N m_i r_i^2)^2} - \frac{c_2^2}{(M - \sum_{i=1}^N m_i r_i^2)^2}\right] - \frac{r_i \dot r_i^2}{1-r_i^2}\\ &+ \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j[r_j(1-r_i^2)\cos(a_i-a_j) - r_i(1-r_i^2)^{\frac12}(1-r_j^2)^{\frac12}\cos(b_i-b_j)]}{(1-\epsilon_{ij}^2)^{\frac32}},\end{aligned} \tag{29}$$

where, for any $i, j \in \{1, 2, \dots, N\}$ *with* $i \neq j$, *we denoted*

$$\epsilon_{ij} := \mathbf{q}_i \cdot \mathbf{q}_j = r_i r_j \cos(a_i - a_j) + (1-r_i^2)^{\frac12}(1-r_j^2)^{\frac12}\cos(b_i - b_j). \tag{30}$$

Proof. Consider a candidate solution of the form (6) for system (3). By expressing each ρ_i in terms of r_i, $i = 1, 2, \dots, N$, we obtain that

$$\rho_i = (1-r_i^2)^{\frac12}, \quad \dot\rho_i = -\frac{r_i\dot r_i}{(1-r_i^2)^{\frac12}}, \quad \ddot\rho_i = -\frac{\dot r_i^2 + r_i(1-r_i^2)\ddot r_i}{(1-r_i^2)^{\frac32}},$$

ϵ_{ij} takes the form (30), and

$$\dot{\mathbf{q}}_i \cdot \dot{\mathbf{q}}_i = \dot r_i^2 + r_i^2\dot\alpha^2 + \frac{r_i^2\dot r_i^2}{1-r_i^2} + (1-r_i^2)\dot\beta^2.$$

Substituting a solution candidate of the form (6) into system (3), and using the above formulas, we obtain for the equations corresponding to $\ddot w_i$ and $\ddot x_i$ the equations

$$\begin{aligned}\ddot r_i &= r_i(1-r_i^2)(\dot\alpha^2 - \dot\beta^2) - \frac{r_i\dot r_i^2}{1-r_i^2}\\ &+ \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j[r_j(1-r_i^2)\cos(a_i-a_j) - r_i(1-r_i^2)^{\frac12}(1-r_j^2)^{\frac12}\cos(b_i-b_j)]}{(1-\epsilon_{ij}^2)^{\frac32}},\end{aligned} \tag{31}$$

$$r_i\ddot\alpha + 2\dot r_i\dot\alpha = -\sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j r_j \sin(a_i - a_j)}{(1-\epsilon_{ij}^2)^{\frac32}}, \quad i = 1, 2, \dots, N, \tag{32}$$

respectively, whereas for the equations corresponding to $\ddot{y}_i, \ddot{z}_i$, we find equations (31) again as well as the equations

$$r_i\ddot{\beta} + 2\dot{r}_i\dot{\beta} = -\sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(1-r_j^2)^{\frac{1}{2}}\sin(b_i - b_j)}{(1-\epsilon_{ij}^2)^{\frac{3}{2}}}, \quad i = 1, 2, \dots, N. \tag{33}$$

We can solve equations (32) the same way we solved equations (15) and obtain

$$\dot{\alpha} = \frac{c_1}{\sum_{i=1}^{N} m_i r_i^2},$$

where c_1 is an integration constant. To solve equations (33), we proceed similarly, with the only change that for each $i = 1, 2, \dots, N$, the corresponding equation gets multiplied by $m_i(1-r_i^2)^{\frac{1}{2}}$ instead of $m_i r_i$, to obtain after addition that

$$\dot{\beta} = \frac{c_2}{M - \sum_{i=1}^{N} m_i r_i^2},$$

where $M = \sum_{i=1}^{N} m_i$ and c_2 is an integration constant. Then equations (31) take the form (29), a remark that completes the proof. □

Remark 8. From (28), we can conclude that $\dot{\alpha}$ and $\dot{\beta}$ are not independent of each other, but connected by the relationship

$$\frac{c_1}{\dot{\alpha}} + \frac{c_2}{\dot{\beta}} = M, \tag{34}$$

written under the assumption that α and β are not constant. In particular, if α and β differ only by an additive constant, they are linear functions of time, i.e.

$$\dot{\alpha} = \dot{\beta} = \frac{c_1 + c_2}{M}. \tag{35}$$

Then system (29) becomes

$$\ddot{r}_i = -\frac{r_i\dot{r}_i^2}{1-r_i^2} + \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j[r_j(1-r_i^2)\cos(a_i - a_j) - r_i(1-r_i^2)^{\frac{1}{2}}(1-r_j^2)^{\frac{1}{2}}\cos(b_i - b_j)]}{(1-\epsilon_{ij}^2)^{\frac{3}{2}}}, \tag{36}$$

$i = 1, 2, \dots, N$.

8.2. **Conservation laws for positive elliptic-elliptic rotopulsators.** In addition to Criterion 2, we would also like to obtain the conservation laws specific to positive elliptic-elliptic rotopulsators. These laws follow by straightforward computations using the above proof, the integral of energy, and the six integrals of the total angular momentum.

Proposition 3. *If system* (3) *has a solution of the form* (6), *then the following expressions are constant:*

— *energy:*

$$h=\sum_{i=1}^N\frac{m_i\dot r_i^2}{2(1-r_i^2)}+\frac{c_1^2}{2\sum_{j=1}^N m_jr_j^2}+\frac{c_2^2}{2(M-\sum_{j=1}^N m_jr_j^2)}-\sum_{1\le i<j\le N}\frac{m_im_j\epsilon_{ij}}{(1-\epsilon_{ij}^2)^{\frac12}};\tag{37}$$

— *total angular momentum relative to the wx-plane,*

$$c_{wx}=c_1,\tag{38}$$

where $c_1\neq 0$ *is the constant in the expression* (28) *of* $\dot\alpha$*;*

— *total angular momentum relative to the wy-plane,*

$$\begin{aligned}c_{wy}=\frac12\sum_{i=1}^N m_i\Big[&r_i(1-r_i^2)^{\frac12}(\dot\alpha+\dot\beta)\sin(\alpha-\beta+a_i-b_i)\\&+r_i(1-r_i^2)^{\frac12}(\dot\alpha-\dot\beta)\sin(\alpha+\beta+a_i+b_i)-\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\cos(\alpha-\beta+a_i-b_i)\\&-\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\cos(\alpha+\beta+a_i+b_i)\Big];\end{aligned}\tag{39}$$

— *total angular momentum relative to the wz-plane,*

$$\begin{aligned}c_{wz}=\frac12\sum_{i=1}^N m_i\Big[&r_i(1-r_i^2)^{\frac12}(\dot\alpha+\dot\beta)\cos(\alpha+\beta+a_i+b_i)\\&+r_i(1-r_i^2)^{\frac12}(\dot\alpha-\dot\beta)\cos(\alpha-\beta+a_i-b_i)+\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\sin(\alpha-\beta+a_i-b_i)\\&-\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\sin(\alpha+\beta+a_i+b_i)\Big];\end{aligned}\tag{40}$$

— *total angular momentum relative to the xy-plane,*

$$\begin{aligned}c_{xy}=-\frac12\sum_{i=1}^N m_i\Big[&r_i(1-r_i^2)^{\frac12}(\dot\alpha+\dot\beta)\cos(\alpha-\beta+a_i-b_i)\\&+r_i(1-r_i^2)^{\frac12}(\dot\alpha-\dot\beta)\cos(\alpha+\beta+a_i+b_i)+\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\sin(\alpha-\beta+a_i-b_i)\\&+\frac{\dot r_i}{(1-r_i^2)^{\frac12}}\sin(\alpha+\beta+a_i+b_i)\Big];\end{aligned}\tag{41}$$

— *total angular momentum relative to the xz-plane,*

$$c_{xz}=\frac{1}{2}\sum_{i=1}^N m_i\Bigg[r_i(1-r_i^2)^{\frac{1}{2}}(\dot\alpha+\dot\beta)\sin(\alpha-\beta+a_i-b_i)$$

$$-r_i(1-r_i^2)^{\frac{1}{2}}(\dot\alpha-\dot\beta)\sin(\alpha+\beta+a_i+b_i)-\frac{\dot r_i}{(1-r_i^2)^{\frac{1}{2}}}\cos(\alpha-\beta+a_i-b_i) \tag{42}$$

$$-\frac{\dot r_i}{(1-r_i^2)^{\frac{1}{2}}}\cos(\alpha+\beta+a_i+b_i)\Bigg];$$

— *total angular momentum relative to the yz-plane,*

$$c_{yz}=c_2, \tag{43}$$

where $c_2\neq 0$ is the constant in the expression (28) *of $\dot\beta$.*

The above result could be also used to prove the nonexistence of some candidates for positive elliptic rotopulsating orbits, by showing that at least one of the above conservation laws is violated.

8.3. **Positive elliptic-elliptic Lagrangian rotopulsators.**

We further provide a class of specific examples of positive elliptic-elliptic rotopulsators of the curved 3-body problem, namely Lagrangian orbits, i.e. bodies that lie at the vertices of a rotating equilateral triangle in $\mathbb{S}^3$. These systems rotate with respect to the wx- and the yz-plane, but have no rotations relative to the other base planes. These solutions provide the first examples of nonstandard relative equilibria, as stated in Definition 6, i.e. every such equilateral triangle remains congruent to itself, in spite of the fact that its orbit cannot be generated by the action of any single element of the underlying Lie group $SO(2)\times SO(2)$, which occurs naturally from the way we fixed the coordinate axes, although it may be generated by an element of $SO(4)$.

Consider three equal masses, $m_1=m_2=m_3=:m$, and a candidate solution of the form

$$\mathbf{q}=(\mathbf{q}_1,\mathbf{q}_2,\mathbf{q}_3),\quad \mathbf{q}_i=(w_i,x_i,y_i,z_i),\quad i=1,2,3, \tag{44}$$

$$w_1=r(t)\cos\alpha(t),\ x_1(t)=r(t)\sin\alpha(t),\ y_1=\rho(t)\cos\beta(t),\ z_1(t)=\rho(t)\sin\beta(t),$$

$$w_2=r(t)\cos[\alpha(t)+2\pi/3],\ x_2(t)=r(t)\sin[\alpha(t)+2\pi/3],$$

$$y_2=\rho(t)\cos[\beta(t)+2\pi/3],\ z_2(t)=\rho(t)\sin[\beta(t)+2\pi/3],$$

$$w_3=r(t)\cos[\alpha(t)+4\pi/3],\ x_3(t)=r(t)\sin[\alpha(t)+4\pi/3],$$

$$y_3=\rho(t)\cos[\beta(t)+4\pi/3],\ z_3(t)=\rho(t)\sin[\beta(t)+4\pi/3],$$

with α and β nonconstant functions and $r^2+\rho^2=1$. We can now prove the following result.

Proposition 4. *Consider the curved* 3*-body problem in* $\mathbb{S}^3$ *given by system* (3)*. Then, except for a negligible set of solutions formed by positive elliptic-elliptic Lagrangian relative equilibria, every candidate solution of the form* (44) *is a positive elliptic-elliptic Lagrangian rotopulsator and, at the same time, a nonstandard relative equilibrium.*

Proof. To prove this result, let us first apply Criterion 2 to a candidate solution of the form (44) for system (3). Straightforward computations show that

$$\epsilon_{12} = \epsilon_{21} = \epsilon_{13} = \epsilon_{31} = \epsilon_{23} = \epsilon_{32} = -1/2,$$

which means that the sides of the equilateral triangle don't vary in time, so if this solution candidate proves to be a positive elliptic-elliptic Lagrangian rotopulsator, then it is also a nonstandard relative equilibrium.

Notice further that

$$\dot\alpha = \frac{c_1}{3mr^2}, \quad \dot\beta = \frac{c_2}{3m(1-r^2)}, \tag{45}$$

and that system (29) reduces to the family of first-order systems

$$\begin{cases} \dot r = u, \\ \dot u = \frac{c_1^2(1-r^2)}{9m^2r^3} - \frac{r(9m^2u^2+c_2^2)}{9m^2(1-r^2)}. \end{cases} \tag{46}$$

As in the proof of Proposition 2, the existence and uniqueness of analytic positive elliptic-elliptic rotopulsators for admissible initial data follows. □

8.4. **The nature of the solutions.** To get some insight into the nature of these solutions, let us first find the fixed points of system (46), which occur for

$$(c_1^2 - c_2^2)r^4 - 2c_1^2r^2 + c_1^2 = 0.$$

For $c_1 \ne c_2$, there are at most two fixed points, $r_1 = \sqrt{\frac{c_1}{c_1-c_2}}$ and $r_2 = \sqrt{\frac{c_1}{c_1+c_2}}$, only one of which is between 0 and 1, whereas for $c_1 = c_2$ there is a single fixed point, namely $r_0 = 1/\sqrt{2}$. So in both cases the fixed point is unique.

Numerical experiments suggest that the phase space picture of system (46) looks like in Fig.(2), with r periodic, a fact in agreement with the Lie theory applied to the group $SO(4)$. Consequently, the positive elliptic-elliptic Lagrangian orbits are expected to be not periodic, since the periods of α and β differ, in general, except for a negligible set corresponding to periodic orbits.

Remark 9. From relations (45) we can see how the angular velocities $\dot\alpha$ and $\dot\beta$ vary relative to r: when r is close to 0, $|\dot\alpha|$ is large, while $|\dot\beta|$ is small, and the other way around when r is close to 1. Moreover, the expressions of $\dot\alpha$ and $\dot\beta$ are consistent with relation (34).



FIGURE 2. The flow of system (46) for $m = 1/3$ and $c_1 = c_2 = 1$. For $c_1 \neq c_2$, the flow looks qualitatively similar.

Remark 10. If we initially assume the masses to be distinct, but the bodies to correspond to the same function r, it follows that the solutions don't exist, which means that the masses must be always equal. If we additionally take distinct functions r_1, r_2, r_3, then

$$\epsilon_{ij} = -\frac{r_i r_j + (1-r_i^2)^{1/2}(1-r_j^2)^{1/2}}{2} = \frac{(r_i - r_j)^2 + [(1-r_i^2)^{1/2} - (1-r_j^2)^{1/2}]^2 - 2}{4},$$

which implies that the triangle is not necessarily equilateral, so the orbit may not be Lagrangian at all.

Remark 11. The energy relation (37) takes the form

$$h = \frac{3m\dot{r}^2}{2(1-r^2)} + \frac{1}{6m^2}\left[\frac{c_1^2}{r^2} + \frac{c_2^2}{1-r^2}\right] + \sqrt{3}m^2,$$

so the energy constant, h, is always positive.

Remark 12. When $\dot{\alpha} = \dot{\beta}$, we have $r^2 = \frac{c_1}{c_1+c_2}$, which means that r must be constant, if it exists, so

$$\dot{\alpha} = \dot{\beta} = \frac{c_1 + c_2}{3m}.$$

These facts imply that the orbit becomes a standard relative equilibrium, not a rotopulsator, and the integrals (37), (39), (40), (41), and (42) are constant, as expected.

It is natural to ask whether there exist other nonstandard relative equilibria beyond the positive elliptic-elliptic Lagrangian rotopulsators described in this section. A close look at expression (30) leads to the following conclusion.

Remark 13. If for for any positive elliptic-elliptic rotopulsator, i.e. any solution of the form (6) of system (3), we have $r_i = r_j$ for all $i, j \in \{1, 2, \dots, N\}$ and $a_i =$

$b_i, i = 1, 2, \dots, N$, then this solution is also a nonstandard relative equilibrium. In particular, positive elliptic-elliptic rotopulsators given by regular polygons can satisfy this property.

9. QUALITATIVE BEHAVIOUR OF THE POSITIVE ROTOPULSATORS

In this section we will prove a result that describes the qualitative behaviour of rotopulsators in $\mathbb{S}^3$. For this purpose, we first briefly introduce an object familiar to geometric topologists and present some of its properties.

9.1. Clifford tori.

The 2-dimensional manifold defined by

$$\mathbf{T}_{r\rho} := \{(w, x, y, z) \in \mathbb{R}^4 \mid r^2 + \rho^2 = 1, \quad 0 \le \theta, \phi < 2\pi\}, \tag{47}$$

where $w = r\cos\theta, x = r\sin\theta, y = \rho\cos\phi$, and $z = \rho\sin\phi$, with $r, \rho \ge 0$, is called a Clifford torus, and it has zero Gaussian curvature. Since the distance from 0 to every point of $\mathbf{T}_{r\rho}$ is 1, it follows that Clifford tori are included in $\mathbb{S}^3$. When r (and, consequently, ρ) takes all the values between 0 and 1, the family of Clifford tori such defined foliates $\mathbb{S}^3$. Each Clifford torus splits $\mathbb{S}^3$ into two solid tori and forms the boundary between them. The two solid tori are congruent only when $r = \rho = 1/\sqrt{2}$.

We have previously shown that positive relative equilibria rotate on Clifford tori, [6], [7]. We will next prove that, at every moment in time, a rotopulsator passes through a different Clifford torus of any given foliation of $\mathbb{S}^3$. In other words, rotopulsators cannot be generated by an element of the underlying Lie group $SO(2) \times SO(2)$.

9.2. Positive rotopulsators.

We can now state and prove the following result.

Theorem 1. *Consider a positive elliptic or positive elliptic-elliptic rotopulsator of the curved N-body in $\mathbb{S}^3$, for which a coordinate system of the ambient space $\mathbb{R}^4$ is fixed. Then, for the corresponding foliation of $\mathbb{S}^3$ given by Clifford tori, the trajectory of each body cannot be contained in any single Clifford torus, but intersects a continuum of Clifford tori. For positive elliptic and positive elliptic-elliptic relative equilibria, however, each body moves on some Clifford torus of the foliation for all time.*

Proof. The qualitative behaviour of positive relative equilibria in $\mathbb{S}^3$, including properties specific to each type of orbit, was previously described and proven to take place, [6], [7], so we will further focus on positive rotopulsators. Since a reference frame was a priori fixed in $\mathbb{R}^4$, the natural foliation of $\mathbb{S}^3$ is given by the family $(\mathbf{T}_{r\rho})_{0\le r,\rho,\le 1}$ of Clifford tori, with $\mathbf{T}_{r\rho}$ defined in (47).

But the positive elliptic-elliptic rotopulsators are given by equations (2), with r and ρ nonconstant functions satisfying the relation $r^2 + \rho^2 = 1$. Each Clifford torus in the foliation, however, corresponds to a constant r and a constant ρ that also satisfy the relation $r^2+\rho^2=1$. Consequently, at every moment in time, each particle passes through a different Clifford torus, thus travelling through a continuum of such objects during its motion.

In the case of positive elliptic rotopulsators, the rotation takes place only with respect to the variables w and x, so we can treat this situation as a particular form of positive elliptic-elliptic rotopulsators, for which the result was proved above. This remark completes the proof. $\square$

Remark 14. The nonstandard relative equilibria (44) behave like the positive elliptic-elliptic Lagrangian rotopulsators, not like the positive elliptic-elliptic Lagrangian relative equilibria, in the sense that they pass through a continuum of Clifford tori instead of moving all along a single Clifford torus, as standard relative equilibria do. But like the relative equilibria, they maintain constant mutual distances during the motion.

10. NEGATIVE ELLIPTIC ROTOPULSATORS

In this section we analyze the solutions given in Definition 3. We first prove a criterion for finding such solutions, then obtain the conservation laws, and finally discuss a particular class of examples, namely the negative elliptic Lagrangian rotopulsators of the 3-body problem in $\mathbb{H}^3$.

10.1. Criterion for negative elliptic rotopulsators.

The following result provides necessary and sufficient conditions for the existence of negative elliptic rotopulsators in $\mathbb{H}^3$.

Criterion 3. *A solution candidate of the form* (7) *is a positive elliptic rotopulsator for system* (4) *if and only if* $c_{yz}=0$,

$$\dot\alpha=\frac{b}{\sum_{j=1}^N m_j(z_j^2-y_j^2-1)}, \tag{48}$$

where b is a nonzero constant, and the variables $y_i, z_i,\ i=1,2,\dots,N$, satisfy the system of $2N$ second-order differential equations

$$\begin{cases}\ddot y_i=\sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(y_j+\mu_{ij}y_i)}{(\mu_{ij}^2-1)^{3/2}}+F_iy_i\\ \ddot z_i=\sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(z_j+\mu_{ij}z_i)}{(\mu_{ij}^2-1)^{3/2}}+F_iz_i,\end{cases} \tag{49}$$

where

$$F_i:=\frac{[(y_i\dot z_i-z_i\dot y_i)^2+\dot z_i^2-\dot y_i^2]}{z_i^2-y_i^2-1}+\frac{b^2(z_i^2-y_i^2-1)}{[\sum_{j=1}^N m_j(z_j^2-y_j^2-1)]^2}, \tag{50}$$

$i=1,2,\dots,N$, and, for any $i,j\in\{1,2,\dots,N\}$, μ_{ij} is given by

$$\mu_{ij}:=\mathbf{q}_i\cdot\mathbf{q_j}=(z_i^2-y_i^2-1)^{\frac12}(z_j^2-y_j^2-1)^{\frac12}\cos(a_i-a_j)+y_iy_j-z_iz_j.$$

Proof. Consider a solution candidate of the form (7) subject to the above initial conditions. Then, for any $i, j \in \{1, 2, \dots, N\}$, we have

$$\mu_{ij} := \mathbf{q}_i \boxdot \mathbf{q_j} = (z_i^2 - y_i^2 - 1)^{\frac{1}{2}}(z_j^2 - y_j^2 - 1)^{\frac{1}{2}} \cos(a_i - a_j) + y_i y_j - z_i z_j,$$

and, for any $i = 1, 2, \dots, N$, we find that

$$\dot{\mathbf{q}}_i \boxdot \dot{\mathbf{q}}_i = \frac{(y_i \dot{z}_i - z_i \dot{y}_i)^2 + \dot{z}_i^2 - \dot{y}_i^2 + (z_i^2 - y_i^2 - 1)^2 \dot{\alpha}^2}{z_i^2 - y_i^2 - 1}.$$

For all $i = 1, 2, \dots, N$, each r_i can be expressed in terms of y_i and z_i to obtain

$$r_i = (z_i^2 - y_i^2 - 1)^{\frac{1}{2}}, \quad \dot{r}_i = \frac{z_i \dot{z}_i - y_i \dot{y}_i}{(z_i^2 - y_i^2 - 1)^{\frac{1}{2}}},$$

$$\ddot{r}_i = \frac{(z_i^2 - y_i^2 - 1)(z_i \ddot{z}_i - y_i \ddot{y}_i) + \dot{y}_i^2 - \dot{z}_i^2 - (y_i \dot{z}_i - z_i \dot{y}_i)^2}{(z_i^2 - y_i^2 - 1)^{\frac{3}{2}}}.$$

Substituting a solution of the form (7) into system (4) and employing the above formulas, we obtain for the equations corresponding to $\ddot{y}_i$ and $\ddot{z}_i$ that

$$\ddot{y}_i = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(y_j + \mu_{ij} y_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(y_i \dot{z}_i - z_i \dot{y}_i)^2 + \dot{z}_i^2 - \dot{y}_i^2] y_i}{z_i^2 - y_i^2 - 1} + (z_i^2 - y_i^2 - 1) y_i \dot{\alpha}^2, \tag{51}$$

$$\ddot{z}_i = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(z_j + \mu_{ij} z_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(y_i \dot{z}_i - z_i \dot{y}_i)^2 + \dot{z}_i^2 - \dot{y}_i^2] z_i}{z_i^2 - y_i^2 - 1} + (z_i^2 - y_i^2 - 1) z_i \dot{\alpha}^2, \tag{52}$$

whereas for the equations corresponding to $\ddot{w}_i$ and $\ddot{x}_i$, after some long computations that also use (51) and (52), we are led either to identities or to the equations

$$r_i \ddot{\alpha} + 2\dot{r}_i \dot{\alpha} = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j r_j \sin(a_j - a_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}}, \quad i = 1, 2, \dots, N. \tag{53}$$

For every $i = 1, 2, \dots, N$, we further multiply the ith equation in (53) by $m_i r_i$, add the resulting N equations, and notice that

$$\sum_{i=1}^{N} \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_i m_j r_i r_j \sin(a_j - a_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}} = 0.$$

Thus we obtain the equation

$$\ddot{\alpha} \sum_{i=1}^{N} m_i r_i^2 + 2\dot{\alpha} \sum_{i=1}^{N} m_i r_i \dot{r}_i = 0,$$

which has the solution

$$\dot\alpha = \frac{b}{\sum_{i=1}^N m_i r_i^2} = \frac{b}{\sum_{i=1}^N m_i(z_i^2 - y_i^2 - 1)},$$

where b is an integration constant. Consequently, equations (51) and (52) become

$$\ddot y_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(y_j + \mu_{ij} y_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(y_i\dot z_i - z_i\dot y_i)^2 + \dot z_i^2 - \dot y_i^2]y_i}{z_i^2 - y_i^2 - 1} + \frac{b^2(z_i^2 - y_i^2 - 1)y_i}{[\sum_{j=1}^N m_j(z_j^2 - y_j^2 - 1)]^2}, \tag{54}$$

$$\ddot z_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(z_j + \mu_{ij} z_i)}{(\mu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(y_i\dot z_i - z_i\dot y_i)^2 + \dot z_i^2 - \dot y_i^2]z_i}{z_i^2 - y_i^2 - 1} + \frac{b^2(z_i^2 - y_i^2 - 1)z_i}{[\sum_{j=1}^N m_j(z_j^2 - y_j^2 - 1)]^2}, \tag{55}$$

$i = 1, 2, \dots, N$. Since equations (54) and (55) are equivalent to system (49), and we have $c_{yz} = 0$ from the definition of negative elliptic rotopulsaating orbits, the criterion is proved. □

Remark 15. Notice that the constant c_{wx} of the total angular momentum is

$$c_{wx} = \sum_{i=1}^N m_i(w_i\dot x_i - \dot w_i x_i) = \dot\alpha \sum_{i=1}^N m_i(z_i^2 - y_i^2 - 1) = b,$$

so b describes the rotation of the particles relative to the wx-plane.

10.2. **Conservation laws for negative elliptic rotopulsating orbits.** In addition to Criterion 3, we would also like to obtain the conservation laws specific to negative elliptic rotopulsating orbits. They follow by straightforward computations using the above proof, the integral of energy, and the six integrals of the total angular momentum.

Proposition 5. *If system* (4) *has a solution of the form* (7), *then the following expressions are constant:*

— *energy,*

$$\begin{aligned} h = &\sum_{i=1}^N \frac{m_i[(y_i\dot z_i - z_i\dot y_i)^2 + \dot z_i^2 - \dot y_i^2]}{2(z_i^2 - y_i^2 - 1)} \\ &+ \frac{b^2}{2\sum_{i=1}^N m_i(z_i^2 - y_i^2 - 1)} + \sum_{1\le i<j\le N} \frac{m_i m_j \mu_{ij}}{(\mu_{ij}^2 - 1)^{\frac{1}{2}}}; \end{aligned} \tag{56}$$

— *total angular momentum relative to the wy-plane,*

$$\begin{aligned} c_{wy} = {} & \sum_{i=1}^{N} m_i\left[(z_i^2-y_i^2-1)^{\frac{1}{2}}\dot{y}_i+\frac{(y_i\dot{y}_i-z_i\dot{z}_i)y_i}{(z_i^2-y_i^2-1)^{\frac{1}{2}}}\right]\cos(\alpha+a_i) \\ & +\frac{b}{\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)}\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)^{\frac{1}{2}}y_i\sin(\alpha+a_i); \end{aligned} \tag{57}$$

— *total angular momentum relative to the wz-plane,*

$$\begin{aligned} c_{wz} = {} & \sum_{i=1}^{N} m_i\left[(z_i^2-y_i^2-1)^{\frac{1}{2}}\dot{z}_i+\frac{(y_i\dot{y}_i-z_i\dot{z}_i)z_i}{(z_i^2-y_i^2-1)^{\frac{1}{2}}}\right]\cos(\alpha+a_i) \\ & +\frac{b}{\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)}\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)^{\frac{1}{2}}z_i\sin(\alpha+a_i); \end{aligned} \tag{58}$$

— *total angular momentum relative to the xy-plane,*

$$\begin{aligned} c_{xy} = {} & \sum_{i=1}^{N} m_i\left[(z_i^2-y_i^2-1)^{\frac{1}{2}}\dot{y}_i+\frac{(y_i\dot{y}_i-z_i\dot{z}_i)y_i}{(z_i^2-y_i^2-1)^{\frac{1}{2}}}\right]\sin(\alpha+a_i) \\ & -\frac{b}{\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)}\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)^{\frac{1}{2}}y_i\cos(\alpha+a_i); \end{aligned} \tag{59}$$

— *total angular momentum relative to the xz-plane,*

$$\begin{aligned} c_{xz} = {} & \sum_{i=1}^{N} m_i\left[(z_i^2-y_i^2-1)^{\frac{1}{2}}\dot{z}_i+\frac{(y_i\dot{y}_i-z_i\dot{z}_i)z_i}{(z_i^2-y_i^2-1)^{\frac{1}{2}}}\right]\sin(\alpha+a_i) \\ & -\frac{b}{\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)}\sum_{i=1}^{N} m_i(z_i^2-y_i^2-1)^{\frac{1}{2}}z_i\cos(\alpha+a_i); \end{aligned} \tag{60}$$

— *total angular momentum relative to the yz-plane,*

$$c_{yz} = \sum_{i=1}^{N} m_i(y_i\dot{z}_i-\dot{y}_iz_i). \tag{61}$$

10.3. **Negative elliptic Lagrangian rotopulsators.**

We further provide a class of specific examples of negative elliptic rotopulsators of the curved 3-body problem, namely Lagrangian orbits in $\mathbb{H}^3$. These systems rotate relative to the wx-plane, but have no rotations relative to other base planes.

Consider three equal masses, $m_1=m_2=m_3=:m$, and a candidate solution of the form

$$\mathbf{q}=(\mathbf{q}_1,\mathbf{q}_2,\mathbf{q}_3),\quad \mathbf{q}_i=(w_i,x_i,y_i,z_i),\quad i=1,2,3, \tag{62}$$

$$
\begin{gathered}
w_1 = r(t)\cos\alpha(t),\ x_1(t) = r(t)\sin\alpha(t),\ y_1 = y(t),\ z_1(t) = z(t),\\
w_2 = r(t)\cos[\alpha(t) + 2\pi/3],\ x_2(t) = r(t)\sin[\alpha(t) + 2\pi/3], y_2 = y(t),\ z_2(t) = z(t),\\
w_3 = r(t)\cos[\alpha(t) + 4\pi/3],\ x_3(t) = r(t)\sin[\alpha(t) + 4\pi/3], y_3 = y(t),\ z_3(t) = z(t).
\end{gathered}
$$

With the help of Criterion 3, we can now show that, in general, these are indeed solutions of system (4).

Proposition 6. *Consider the curved* 3*-body problem in* $\mathbb{H}^3$ *given by system* (4)*. Then, except for a negligible set of solutions formed by negative elliptic Lagrangian relative equilibria, every candidate solution of the form* (62) *is a negative elliptic Lagrangian rotopulsator.*

Proof. Let us consider a candidate solution of the form (62). Then, using Criterion 3, straightforward computations show that

$$
\mu_{12} = \mu_{13} = \mu_{23} = \frac{3y^2 - 3z^2 + 1}{2}, \quad \dot\alpha = \frac{b}{3m(z^2 - y^2 - 1)}
$$

and that the variables y and z satisfy the equations

$$
\begin{cases}
\ddot y = G(y, z, \dot y, \dot z)y\\
\ddot z = G(y, z, \dot y, \dot z)z,
\end{cases}
\tag{63}
$$

where

$$
\begin{aligned}
G(y, z, \dot y, \dot z) = {} & \frac{(y\dot z - \dot y z)^2 + \dot z^2 - \dot y^2}{z^2 - y^2 - 1} + \frac{b^2}{9m^2(z^2 - y^2 - 1)}\\
& - \frac{8m}{\sqrt{3}(z^2 - y^2 - 1)^{\frac{1}{2}}(3z^2 - 3y^2 + 1)^{\frac{3}{2}}}.
\end{aligned}
$$

From (63), we can conclude that $\ddot y z = y\ddot z$, which implies that

$$
y\dot z - z\dot y = k \text{ (constant)}.
$$

But, since from (61) we have that $3m(y\dot z - z\dot y) = c_{yz}$, it follows that $k = c_{yz}/3m$.

Notice that the energy relation (56) takes the form

$$
\frac{3m[(y\dot z - \dot y z)^2 + \dot z^2 - \dot y^2]}{2(z^2 - y^2 - 1)} + \frac{b^2}{6m(z^2 - y^2 - 1)} + \frac{\sqrt{3}m^2(3y^2 - 3z^2 + 1)}{(z^2 - y^2 - 1)^{\frac{1}{2}}(3z^2 - 3y^2 + 1)^{\frac{1}{2}}} = h,
$$

which implies that G can be written as

$$
G(y, z) = \frac{2h}{3m} - \frac{2m[5 - 9(y^2 - z^2)^2]}{\sqrt{3}(z^2 - y^2 - 1)^{\frac{1}{2}}(3z^2 - 3y^2 + 1)^{\frac{3}{2}}}.
$$

Since

$$
\sin\alpha + \sin(\alpha + 2\pi/3) + \sin(\alpha + 4\pi/3) = \cos\alpha + \cos(\alpha + 2\pi/3) + \cos(\alpha + 4\pi/3) = 0,
$$

it follows from (57), (58), (59), and (60) that $c_{wy} = c_{wz} = c_{xy} = c_{xz} = 0$, so the triangle has no rotation relative to the planes wy, wz, xy, and xz. Since there is no rotation relative to the plane yz either, i.e. $c_{yz} = 0$, we have $k = 0$, so $y\dot{z} - z\dot{y} = 0$, and if we assume that z does not take zero values, we can conclude that $\frac{d}{dt}\frac{y}{z} = 0$, so $y(t) = \gamma z(t)$, where γ is a constant. If we denote $\epsilon = 1 - \gamma^2$, system (63) reduces to the family of first-order systems

$$\begin{cases} \dot{z} = u \\ \dot{u} = \left[\frac{2h}{3m} - \frac{2m(5-9\epsilon^2 z^4)}{\sqrt{3}(\epsilon z^2 - 1)^{\frac{1}{2}}(3\epsilon z^2+1)^{\frac{3}{2}}}\right] z. \end{cases} \tag{64}$$

Notice that since a point (w, x, y, z) on $\mathbb{H}^3$ satisfies the equation $w^2 + x^2 + y^2 - z^2 = -1$ and $z \geq 1$, we necessarily have that $\epsilon \geq 1$.

Standard results of the theory of ordinary differential equations can now be applied to system (64) to prove the existence and uniqueness of analytic negative elliptic Lagrangian rotopulsators, for admissible initial conditions. To show that, except for a negligible set, they are all rotopulsators, we identify the relative equilibria, which are fixed points of system (64).

One fixed point of the vector field in (64) is obviously $(z, u) = (0, 0)$, but it lies outside the domain $z \geq 1$. The other fixed points, if any, must be of the form $(z, 0)$, where the positive values of z are given by the roots of the polynomial

$$Q(z) = 27(h^2 - 9m^4)\epsilon^4 z^8 - 18(h^2 - 15m^4)\epsilon^2 z^4 - 8h^2\epsilon z^2 - h^2 - 75m^4.$$

By Descartes's rule of signs, we must distinguish between two cases:

(i) $|h| < \sqrt{15}m^2$, when Q has no positive roots at all;

(ii) $|h| \geq \sqrt{15}m^2$, when Q has exactly one positive root.

Case (i) leads to no fixed points for system (64). Case (ii) always leads to one fixed point since the unique positive root, $z_0 := z_0(m, h, \epsilon)$, is also larger than 1. To show this fact, let $\xi = \epsilon z^2$. Then we can associate to Q the polynomial

$$\bar{Q}(\xi) = 27(h^2 - 9m^4)\xi^4 - 18(h^2 - 15m^4)\xi^2 - 8h^2\xi - h^2 - 75m^4.$$

Since $\epsilon \geq 1$, to prove that $z_0 \geq 1$ it is enough show that $\xi_0 := \epsilon z_0 \geq 1$. For this, notice that $\bar{Q}(1) = -48m^4 < 0$. Since $\lim_{\xi \to +\infty} \bar{Q}(\xi) = +\infty$, $\bar{Q}$ has a single positive root that is also larger than 1. This remark completes the proof. $\square$

11. NEGATIVE HYPERBOLIC ROTOPULSATORS

In this section we analyze the solutions given in Definition 4. We first prove a criterion for finding such solutions, then provide the conservation laws, and finally discuss a particular class of examples, namely the negative hyperbolic Eulerian rotopulsators of the 3-body problem in $\mathbb{H}^3$.

11.1. **Criterion for negative hyperbolic rotopulsators.** The following result provides necessary and sufficient conditions for the existence of negative hyperbolic rotopulsators in $\mathbb{H}^3$.

Criterion 4. *A solution candidate of the form* (8) *is a negative hyperbolic rotopulsating orbit for system* (4) *if and only if* $c_{wx} = 0$,

$$\dot\beta = \frac{a}{\sum_{j=1}^N m_j(w_j^2 + x_j^2 + 1)}, \tag{65}$$

where a *is a constant, and the variables* $w_i, x_i,\ i = 1, 2, \dots, N$, *satisfy the system of* $2N$ *second-order differential equations*

$$\begin{cases} \ddot w_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(w_j+\nu_{ij}w_i)}{(\nu_{ij}^2-1)^{3/2}} + H_i w_i \\ \ddot x_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(x_j+\nu_{ij}x_i)}{(\nu_{ij}^2-1)^{3/2}} + H_i x_i, \end{cases} \tag{66}$$

where

$$H_i := \frac{(w_i\dot x_i - x_i\dot w_i)^2 + \dot w_i^2 + \dot x_i^2}{w_i^2 + x_i^2 + 1} + \frac{a^2(w_i^2 + x_i^2 + 1)}{[\sum_{j=1}^N m_j(w_j^2 + x_j^2 + 1)]^2}, \tag{67}$$

$i = 1, 2, \dots, N$, *and, for any* $i, j \in \{1, 2, \dots, N\}$, ν_{ij} *is given by*

$$\nu_{ij} := \mathbf{q}_i \boxdot \mathbf{q_j} = w_i w_j + x_i x_j - (w_i^2 + x_i^2 + 1)^{\frac12}(w_j^2 + x_j^2 + 1)^{\frac12}\cosh(b_i - b_j). \tag{68}$$

Proof. Consider a solution candidate of the form (8) subject to the above initial conditions. Then, for any $i, j \in \{1, 2, \dots, N\}$ we have

$$\nu_{ij} := \mathbf{q}_i \boxdot \mathbf{q_j} = w_i w_j + x_i x_j - (w_i^2 + x_i^2 + 1)^{\frac12}(w_j^2 + x_j^2 + 1)^{\frac12}\cosh(b_i - b_j),$$

and for any $i = 1, 2, \dots, N$ we find that

$$\dot{\mathbf{q}}_i \boxdot \dot{\mathbf{q}}_i = \frac{(w_i\dot x_i - x_i\dot w_i)^2 + \dot w_i^2 + \dot x_i^2 + (w_i^2 + x_i^2 + 1)^2\dot\beta^2}{w_i^2 + x_i^2 + 1}.$$

For all $i = 1, 2, \dots, N$, each r_i can be expressed in terms of y_i and z_i to obtain

$$\eta_i = (w_i^2 + x_i^2 + 1)^{\frac12}, \quad \dot\eta_i = \frac{w_i\dot w_i + x_i\dot x_i}{(w_i^2 + x_i^2 + 1)^{\frac12}},$$

$$\ddot\eta_i = \frac{(w_i^2 + x_i^2 + 1)(w_i\ddot w_i + x_i\ddot x_i) + \dot w_i^2 + \dot x_i^2 + (w_i\dot x_i - x_i\dot w_i)^2}{(w_i^2 + x_i^2 + 1)^{\frac32}}.$$

Substituting a solution of the form (8) into system (4) and employing the above formulas, we obtain for the equations corresponding to $\ddot w_i$ and $\ddot x_i$ that

$$\ddot w_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(w_j + \nu_{ij}w_i)}{(\nu_{ij}^2 - 1)^{\frac32}} + \frac{[(w_i\dot x_i - x_i\dot w_i)^2 + \dot w_i^2 + \dot x_i^2]w_i}{w_i^2 + x_i^2 + 1} + (w_i^2 + x_i^2 + 1)w_i\dot\beta^2, \tag{69}$$

$$\ddot x_i = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j(x_j + \nu_{ij}x_i)}{(\mu_{ij}^2 - 1)^{\frac32}} + \frac{[(w_i\dot x_i - x_i\dot w_i)^2 + \dot w_i^2 + \dot x_i^2]x_i}{w_i^2 + x_i^2 + 1} + (w_i^2 + x_i^2 + 1)x_i\dot\beta^2, \tag{70}$$

whereas for the equations corresponding to $\ddot{y}_i$ and $\ddot{z}_i$, after some long computations that also use (69) and (70), we are led either to identities or to the equations

$$\eta_i\ddot{\beta} + 2\dot{\eta}_i\dot{\beta} = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j\eta_j \sinh(b_j - b_i)}{(\nu_{ij}^2 - 1)^{\frac{3}{2}}}, \quad i = 1, 2, \dots, N. \tag{71}$$

For every $i = 1, 2, \dots, N$, we further multiply the ith equation in (53) by $m_i r_i$, add the resulting N equations, and notice that

$$\sum_{i=1}^{N}\sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_i m_j \eta_i\eta_j \sinh(b_j - b_i)}{(\nu_{ij}^2 - 1)^{\frac{3}{2}}} = 0.$$

Thus we obtain the equation

$$\Big(\sum_{i=1}^{N} m_i\eta_i^2\Big)\ddot{\beta} + 2\Big(\sum_{i=1}^{N} m_i\eta_i\dot{\eta}_i\Big)\dot{\beta} = 0,$$

which has the solution

$$\dot{\beta} = \frac{a}{\sum_{i=1}^{N} m_i\eta_i^2} = \frac{a}{\sum_{i=1}^{N} m_i(w_i^2 + x_i^2 + 1)},$$

where a is an integration constant. Consequently, equations (69) and (70) become

$$\ddot{w}_i = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(w_j + \nu_{ij}w_i)}{(\nu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(w_i\dot{x}_i - x_i\dot{w}_i)^2 + \dot{w}_i^2 + \dot{x}_i^2]w_i}{w_i^2 + x_i^2 + 1} + \frac{a^2(w_i^2 + x_i^2 + 1)w_i}{[\sum_{j=1}^{N} m_j(w_j^2 + x_j^2 + 1)]^2}, \tag{72}$$

$$\ddot{x}_i = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(x_j + \nu_{ij}x_i)}{(\nu_{ij}^2 - 1)^{\frac{3}{2}}} + \frac{[(w_i\dot{x}_i - x_i\dot{w}_i)^2 + \dot{w}_i^2 + \dot{x}_i^2]x_i}{w_i^2 + x_i^2 + 1} + \frac{a^2(w_i^2 + x_i^2 + 1)x_i}{[\sum_{j=1}^{N} m_j(w_j^2 + x_j^2 + 1)]^2}, \tag{73}$$

$i = 1, 2, \dots, N$. Since equations (72) and (73) lead to system (66), and the condition $c_{wx} = 0$ occurs from Definition 4, the criterion is proved. □

Remark 16. Notice that the constant c_{yz} of the angular momentum is

$$c_{yz} = \sum_{i=1}^{N} m_i(y_i\dot{z}_i - \dot{y}_i z_i) = -\dot{\beta}\sum_{i=1}^{N} m_i(w_i^2 + x_i^2 + 1) = -a.$$

11.2. **Conservation laws for negative hyperbolic rotopulsating orbits.** In addition to Criterion 4, we would also like to obtain the conservation laws specific to negative elliptic rotopulsating orbits. They follow by straightforward computations using the above proof, the integral of energy, and the six integrals of the total angular momentum.

Proposition 7. *If system* (4) *has a solution of the form* (8), *then the following expressions are constant:*

— *energy,*

$$\begin{aligned} h = \sum_{i=1}^N \frac{m_i[(w_i\dot{x}_i - x_i\dot{w}_i)^2 + \dot{w}_i^2 + \dot{x}_i^2]}{2(w_i^2+x_i^2+1)} \\ + \frac{a^2}{2\sum_{j=1}^N m_j(w_j^2+x_j^2+1)} + \sum_{1\le i<j\le N} \frac{m_im_j\nu_{ij}}{(\nu_{ij}^2-1)^{1/2}}; \end{aligned} \tag{74}$$

— *total angular momentum relative to the wx-plane,*

$$c_{wx} = \sum_{i=1}^N m_i(w_i\dot{x}_i - \dot{w}_ix_i); \tag{75}$$

— *total angular momentum relative to the wy-plane,*

$$\begin{aligned} c_{wy} = \sum_{i=1}^N \frac{m_i[x_i(w_i\dot{x}_i - x_i\dot{w}_i) - \dot{w}_i]}{(w_i^2+x_i^2+1)^{\frac{1}{2}}}\sinh(\beta+b_i) \\ + \frac{a}{\sum_{j=1}^N m_j(w_j^2+x_j^2+1)} \sum_{i=1}^N m_iw_i(w_i^2+x_i^2+1)^{\frac{1}{2}}\cosh(\beta+b_i); \end{aligned} \tag{76}$$

— *total angular momentum relative to the wz-plane,*

$$\begin{aligned} c_{wz} = \sum_{i=1}^N \frac{m_i[x_i(w_i\dot{x}_i - x_i\dot{w}_i) - \dot{w}_i]}{(w_i^2+x_i^2+1)^{\frac{1}{2}}}\cosh(\beta+b_i) \\ + \frac{a}{\sum_{j=1}^N m_j(w_j^2+x_j^2+1)} \sum_{i=1}^N m_iw_i(w_i^2+x_i^2+1)^{\frac{1}{2}}\sinh(\beta+b_i); \end{aligned} \tag{77}$$

— *total angular momentum relative to the xy-plane,*

$$\begin{aligned} c_{xy} = \sum_{i=1}^N \frac{m_i[w_i(\dot{w}_ix_i - w_i\dot{x}_i) - \dot{x}_i]}{(w_i^2+x_i^2+1)^{\frac{1}{2}}}\sinh(\beta+b_i) \\ + \frac{a}{\sum_{j=1}^N m_j(w_j^2+x_j^2+1)} \sum_{i=1}^N m_ix_i(w_i^2+x_i^2+1)^{\frac{1}{2}}\cosh(\beta+b_i); \end{aligned} \tag{78}$$

— *total angular momentum relative to the xz-plane,*

$$\begin{aligned} c_{xz} = \sum_{i=1}^{N} & \frac{m_i[w_i(\dot{w}_i x_i - w_i \dot{x}_i) - \dot{x}_i]}{(w_i^2 + x_i^2 + 1)^{\frac{1}{2}}} \cosh(\beta + b_i) \\ & + \frac{a}{\sum_{j=1}^{N} m_j (w_j^2 + x_j^2 + 1)} \sum_{i=1}^{N} m_i x_i (w_i^2 + x_i^2 + 1)^{\frac{1}{2}} \sinh(\beta + b_i). \end{aligned} \tag{79}$$

11.3. **Negative hyperbolic Eulerian rotopulsators.** We further provide a class of specific negative hyperbolic rotopulsators of the curved 3-body problem, namely Eulerian orbits in $\mathbb{H}^3$. These systems rotate relative to the yz-plane, but have no rotations relative to other base planes.

Consider three equal masses, $m_1 = m_2 = m_3 =: m$, and a candidate solution of the form

$$\begin{gathered} \mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, 3, \\ w_1 = 0, \quad x_1 = 0, \quad y_1 = \sinh\beta(t), \quad z_1 = \cosh\beta(t), \\ w_1 = w(t), \quad x_1 = x(t), \quad y_1 = \eta(t)\sinh\beta(t), \quad z_1 = \eta(t)\cosh\beta(t), \\ w_1 = -w(t), \quad x_1 = -x(t), \quad y_1 = \eta(t)\sinh\beta(t), \quad z_1 = \eta(t)\cosh\beta(t), \end{gathered} \tag{80}$$

with $w^2 + x^2 - \eta^2 = -1$. With the help of Criterion 4, we can now show that, in general, these are indeed solutions of system (4).

Proposition 8. *Consider the curved* 3*-body problem in* $\mathbb{H}^3$ *given by system* (4)*. Then, except for a negligible set of solutions formed by negative hyperbolic Eulerian relative equilibria, every candidate solution of the form* (80) *is a negative hyperbolic Eulerian rotopulsator.*

Proof. Let us consider a candidate solution of the form (80). Then, using Criterion 4, straightforward computations show that

$$\nu_{12} = \nu_{13} = (w^2 + x^2)^{\frac{1}{2}}, \quad \nu_{23} = -2(w^2 + x^2), \quad \dot{\beta} = \frac{a}{m(2w^2 + 2x^2 + 3)},$$

and the equations of motion reduce to the system

$$\begin{cases} \ddot{w} = J(w, x, \dot{w}, \dot{x})w \\ \ddot{x} = J(w, x, \dot{w}, \dot{x})x, \end{cases} \tag{81}$$

where

$$\begin{aligned} J(w, x, \dot{w}, \dot{x}) = & \frac{m(w^2 + x^2)^{\frac{1}{2}}}{(w^2 + x^2 - 1)^{\frac{3}{2}}} - \frac{m}{(2w^2 + 2x^2 + 1)^{\frac{1}{2}}(2w^2 + 2x^2 - 1)^{\frac{3}{2}}} \\ & + \frac{(w\dot{x} - x\dot{w})^2 + \dot{w}^2 + \dot{x}^2}{w^2 + x^2 + 1} + \frac{a^2(w^2 + x^2 + 1)}{m^2(2w^2 + 2x^2 + 3)^2}. \end{aligned}$$

Notice that the energy integral (74) takes the form

$$\frac{m[(w\dot{x}-x\dot{w})^2+\dot{w}^2+\dot{x}^2]}{w^2+x^2+1}+\frac{a^2}{2m(2w^2+2x^2+3)}$$

$$+\frac{2m^2(w^2+x^2)^{\frac{1}{2}}}{(w^2+x^2-1)^{\frac{1}{2}}}-\frac{2m^2(w^2+x^2)}{(2w^2+2x^2+1)^{\frac{1}{2}}(2w^2+2x^2-1)^{\frac{1}{2}}}=h,$$

which implies that J can be written as

$$J(w,x)=\frac{h}{m}+\frac{m[2(w^2+x^2)(2w^2+2x^2-1)+1]}{(2w^2+2x^2+1)^{\frac{1}{2}}(2w^2+2x^2-1)^{\frac{3}{2}}}$$

$$-\frac{m(w^2+x^2)^{\frac{1}{2}}(2w^2+2x^2-3)}{(w^2+x^2-1)^{\frac{3}{2}}}-\frac{a^2}{2m^2(2w^2+2x^2+3)^2}.$$

From (81), we can conclude that $\ddot{w}x=w\ddot{x}$, which implies that

$$w\dot{x}-\dot{w}x=k \text{ (constant)}.$$

But since from (75) we have that $2m(w\dot{x}-\dot{w}x)=c_{wx}$, it follows that $k=\frac{c_{wx}}{2m}$. From (76), (77), (78), and (79), we can conclude that $c_{wy}=c_{wz}=c_{xy}=c_{xz}=0$, so the orbit has no rotation relative to the planes wy, wz, xy, and xz. Since there is no rotation relative to the plane wx either, i.e. $c_{wx}=0$, we have $k=0$, so $w\dot{x}-\dot{w}x=0$, and if we assume that x does not take zero values, we can conclude that $\frac{d}{dt}\frac{w}{x}=0$, so $w(t)=\xi x(t)$, where ξ is a constant. If we denote $\zeta=\xi^2+1\geq 1$, equations (81) reduce to the family of first-order systems

$$\text{(82)}\qquad \begin{cases}\dot{x}=u\\ \dot{u}=\left[\frac{h}{m}+\frac{m(4\zeta^2x^4-2\zeta x^2+1)}{(2\zeta x^2+1)^{1/2}(2\zeta x^2-1)^{3/2}}-\frac{m(\zeta x^2)^{1/2}(2\zeta x^2-3)}{(\zeta x^2-1)^{3/2}}-\frac{a^2}{2m^2(2\zeta x^2+3)^2}\right]x.\end{cases}$$

Standard results of the theory of differential equations can now be applied to system (82) to prove the existence and uniqueness of analytic negative hyperbolic Eulerian rotopulsators, for admissible initial conditions. The relative equilibria that occur from fixed points of system (82) are given by polynomial equations, so they form a negligible set. This remark completes the proof. □

12. NEGATIVE ELLIPTIC-HYPERBOLIC ROTOPULSATORS

In this section we analyze the solutions given in Definition 5. We first prove a criterion for finding such solutions, then provide the conservation laws, and finally discuss a particular class of examples, namely the negative elliptic-hyperbolic Eulerian rotopulsators of the 3-body problem in $\mathbb{H}^3$.

12.1. **Criterion for negative elliptic-hyperbolic rotopulsators.** The following result provides necessary and sufficient conditions for the existence of negative elliptic-hyperbolic rotopulsators in $\mathbb{H}^3$.

Criterion 5. *A solution candidate of the form* (9) *is a negative elliptic-hyperbolic rotopulsator for system* (4) *if and only if*

$$\dot\alpha = \frac{d_1}{\sum_{i=1}^N m_i r_i^2}, \quad \dot\beta = \frac{d_2}{M + \sum_{i=1}^N m_i r_i^2}, \tag{83}$$

with d_1, d_2 nonzero constants, and the variables r_i, $i = 1, \dots, N$, satisfy the N second-order differential equations

$$\begin{aligned} \ddot r_i = {} & r_i(1+r_i^2)\left[\frac{d_1^2}{(\sum_{i=1}^N m_i r_i^2)^2} - \frac{d_2^2}{(M+\sum_{i=1}^N m_i r_i^2)^2}\right] + \frac{r_i \dot r_i^2}{1+r_i^2} \\ & + \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j[r_j(1+r_i^2)\cos(a_i-a_j) - r_i(1+r_i^2)^{\frac12}(1+r_j^2)^{\frac12}\cosh(b_i-b_j)]}{(\delta_{ij}^2-1)^{\frac32}}, \end{aligned} \tag{84}$$

where, for any $i, j \in \{1, 2, \dots, N\}$ with $i \neq j$, we denoted

$$\delta_{ij} := \mathbf{q}_i \boxdot \mathbf{q}_j = r_i r_j \cos(a_i - a_j) - (1+r_i^2)^{\frac12}(1+r_j^2)^{\frac12}\cosh(b_i - b_j).$$

Proof. Consider a candidate solution of the form (9) for system (4). By expressing each ρ_i in terms of r_i, $i = 1, 2, \dots, N$, we obtain that

$$\eta_i = (1+r_i^2)^{\frac12}, \quad \dot\eta_i = \frac{r_i \dot r_i}{(1+r_i^2)^{\frac12}}, \quad \ddot\eta_i = \frac{\dot r_i^2 + r_i(1+r_i^2)\ddot r_i}{(1+r_i^2)^{\frac32}},$$

$$\delta_{ij} := \mathbf{q}_i \boxdot \mathbf{q}_j = r_i r_j \cos(a_i - a_j) - (1+r_i^2)^{\frac12}(1+r_j^2)^{\frac12}\cosh(b_i - b_j),$$

$$\dot{\mathbf{q}}_i \boxdot \dot{\mathbf{q}}_i = \dot r_i^2 + r_i^2\dot\alpha^2 - \frac{r_i^2 \dot r_i^2}{1+r_i^2} - (1+r_i^2)\dot\beta^2.$$

Substituting a solution candidate of the form (9) into system (4), and using the above formulas, we obtain for the equations corresponding to $\ddot w_i$ and $\ddot x_i$ the equations

$$\begin{aligned} \ddot r_i = {} & r_i(1+r_i^2)(\dot\alpha^2 - \dot\beta^2) + \frac{r_i \dot r_i^2}{1-r_i^2} \\ & + \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j[r_j(1+r_i^2)\cos(a_i-a_j) - r_i(1+r_i^2)^{\frac12}(1+r_j^2)^{\frac12}\cosh(b_i-b_j)]}{(\delta_{ij}^2-1)^{\frac32}}, \end{aligned} \tag{85}$$

$$r_i\ddot\alpha + 2\dot r_i\dot\alpha = \sum_{\substack{j=1\\ j\neq i}}^N \frac{m_j r_j \sin(a_j - a_i)}{(\delta_{ij}^2-1)^{\frac32}}, \quad i = 1, 2, \dots, N, \tag{86}$$

whereas for the equations corresponding to $\ddot{y}_i, \ddot{z}_i$, we find equations (85) again as well as the equations

$$r_i\ddot{\beta} + 2\dot{r}_i\dot{\beta} = \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{m_j(1+r_j^2)^{\frac{1}{2}}\sin(b_j - b_i)}{(\delta_{ij}^2 - 1)^{\frac{3}{2}}}, \quad i = 1, 2, \dots, N. \tag{87}$$

We can solve equations (86) the same way we solved equations (15) and obtain

$$\dot{\alpha} = \frac{d_1}{\sum_{i=1}^{N} m_i r_i^2},$$

where d_1 is an integration constant. To solve equations (87), we proceed similarly, with the only change that for each $i = 1, 2, \dots, N$, the corresponding equation gets multiplied by $m_i(1+r_i^2)^{\frac{1}{2}}$ instead of $m_i r_i$, to obtain after addition that

$$\dot{\beta} = \frac{d_2}{M + \sum_{i=1}^{N} m_i r_i^2},$$

where $M = \sum_{i=1}^{N} m_i$ and d_2 is an integration constant. Then equations (85) take the form (84), a remark that completes the proof. □

Remark 17. Notice that the constants c_{wx} and c_{yz} of the total angular momentum are

$$c_{wx} = \sum_{i=1}^{N} m_i(w_i\dot{x}_i - \dot{w}_i x_i) = \dot{\alpha}\sum_{i=1}^{N} m_i r_i^2 = d_1,$$

$$c_{yz} = \sum_{i=1}^{N} m_i(y_i\dot{z}_i - \dot{z}_i y_i) = -\dot{\beta}\sum_{i=1}^{N} m_i(1 + r_i^2) = -d_2,$$

respectively, so d_1 and d_2 describe the rotation of the particle system relative to the wx-plane and yz-plane, respectively.

Remark 18. From (83), we can conclude that $\dot{\alpha}$ and $\dot{\beta}$ are not independent of each other, but connected by the relationship

$$\frac{d_2}{\dot{\beta}} - \frac{d_1}{\dot{\alpha}} = M. \tag{88}$$

The following observation refers to all the five criteria proved in this paper.

Remark 19. There are 2 first-order and $2N$ second-order differential equations in Criteria 2 and 5, but only 1 first-order and N second-order differential equations in Criteria 1, 3, and 4. This fact, however, leads to no contradiction because the position and velocity of each body m_i in $\mathbb{S}^3$ or $\mathbb{H}^3$ is given by 6 scalar variables, namely $(\alpha, \beta, r_i, \dot{\alpha}, \dot{\beta}, \dot{r}_i)$ for Criteria 2 and 5, $(\beta, w_i, x_i, \dot{\beta}, \dot{w}_i, \dot{x}_i)$ for Criterion 4, and $(\alpha, y_i, z_i, \dot{\alpha}, \dot{y}_i, \dot{z}_i)$ for Criteria 1 and 3, in agreement with the statements of these results.

12.2. **Conservation laws for negative elliptic-hyperbolic rotopulsators.** In addition to Criterion 5, we would also like to obtain the conservation laws specific to negative elliptic rotopulsating orbits. They follow by straightforward computations using the above proof, the integral of energy, and the six integrals of the total angular momentum.

Proposition 9. *If system* (4) *has a solution of the form* (9)*, then the following expressions are constant:*

— *energy,*

$$h=\sum_{i=1}^N\frac{m_i\dot r_i^2}{2(1+r_i^2)}+\frac{d_1^2}{2\sum_{j=1}^N m_jr_j^2}+\frac{d_2^2}{2(M+\sum_{j=1}^N m_jr_j^2)}+\sum_{1\le i<j\le N}\frac{m_im_j\delta_{ij}}{(\delta_{ij}^2-1)^{\frac12}}; \tag{89}$$

— *total angular momentum relative to the wx-plane,*

$$\begin{aligned} c_{wy}=&-\sum_{i=1}^N\frac{m_i\dot r_i}{(1+r_i^2)^{\frac12}}\cos(\alpha+a_i)\sinh(\beta+b_i)\\ &+\sum_{i=1}^N m_ir_i(1+r_i^2)^{\frac12}[\dot\alpha\sin(\alpha+a_i)\sinh(\beta+b_i)+\dot\beta\cos(\alpha+a_i)\cosh(\beta+b_i)]; \end{aligned} \tag{90}$$

— *total angular momentum relative to the wz-plane,*

$$\begin{aligned} c_{wz}=&-\sum_{i=1}^N\frac{m_i\dot r_i}{(1+r_i^2)^{\frac12}}\cos(\alpha+a_i)\cosh(\beta+b_i)\\ &+\sum_{i=1}^N m_ir_i(1+r_i^2)^{\frac12}[\dot\alpha\sin(\alpha+a_i)\cosh(\beta+b_i)+\dot\beta\cos(\alpha+a_i)\sinh(\beta+b_i)]; \end{aligned} \tag{91}$$

— *total angular momentum relative to the xy-plane,*

$$\begin{aligned} c_{xy}=&-\sum_{i=1}^N\frac{m_i\dot r_i}{(1+r_i^2)^{\frac12}}\sin(\alpha+a_i)\sinh(\beta+b_i)\\ &+\sum_{i=1}^N m_ir_i(1+r_i^2)^{\frac12}[\dot\beta\sin(\alpha+a_i)\cosh(\beta+b_i)-\dot\alpha\cos(\alpha+a_i)\sinh(\beta+b_i)]; \end{aligned} \tag{92}$$

— *total angular momentum relative to the xz-plane,*

$$\begin{aligned} c_{xz}=&-\sum_{i=1}^N\frac{m_i\dot r_i}{(1+r_i^2)^{\frac12}}\sin(\alpha+a_i)\cosh(\beta+b_i)\\ &+\sum_{i=1}^N m_ir_i(1+r_i^2)^{\frac12}[\dot\beta\sin(\alpha+a_i)\sinh(\beta+b_i)-\dot\alpha\cos(\alpha+a_i)\cosh(\beta+b_i)]. \end{aligned} \tag{93}$$

12.3. **Negative elliptic-hyperbolic Eulerian rotopulsators.** We further provide a class of specific negative elliptic-hyperbolic rotopulsators of the curved 3-body problem in $\mathbb{H}^3$, namely Eulerian orbits. These systems rotate relative to the wx and yz planes, but have no rotation with respect to the other base planes.

Consider three equal masses, $m_1 = m_2 = m_3 := m$, and a candidate solution of the form

$$\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3), \quad \mathbf{q}_i = (w_i, x_i, y_i, z_i), \quad i = 1, 2, 3, \tag{94}$$

$$w_1 = 0,\ x_1 = 0,\ y_1 = \sinh\beta(t),\ z_1(t) = \cosh\beta(t),$$

$$w_2 = r(t)\cos\alpha(t),\ x_2 = r(t)\sin\alpha(t),\ y_2 = \eta(t)\sinh\beta(t),\ z_2(t) = \eta(t)\cosh\beta(t),$$

$$w_3 = -r(t)\cos\alpha(t),\ x_3 = -r(t)\sin\alpha(t),\ y_3 = \eta(t)\sinh\beta(t),\ z_3 = \eta(t)\cosh\beta(t),$$

with α and β nonconstant functions and $r^2 - \eta^2 = -1$. We can now prove a result which shows that, in general, these orbits exist.

Proposition 10. *Consider the curved* 3*-body problem in* $\mathbb{H}^3$ *given by system* (4)*. Then, except for a negligible set of solutions formed by the negative elliptic-hyperbolic Eulerian relative equilibria, every candidate solution of the form* (94) *is a negative elliptic-hyperbolic Eulerian rotopulsator, which rotates only relative to the planes* wx *and* yz*.*

Proof. Consider a candidate solution of the form (94) of system (4). Then the variables relevant to Criterion 5 take the form

$$\delta_{12} = \delta_{21} = \delta_{13} = \delta_{31} = -(1 + r^2)^{1/2}, \quad \delta_{23} = \delta_{32} = -2r^2 - 1,$$

$$\dot\alpha = \frac{d_1}{2mr^2}, \quad \dot\beta = \frac{d_2}{m(3 + 2r^2)}, \quad \text{with } d_1, d_2 \neq 0 \text{ constants.} \tag{95}$$

In system (84), the equation corresponding to $\ddot r_1$ is an identity, whereas the equations corresponding to $\ddot r_2$ and $\ddot r_3$ are identical, given by

$$\ddot r = r(1 + r^2)(\dot\alpha^2 - \dot\beta^2) + \frac{r\dot r^2}{1 + r^2} - \frac{m(5 + 4r^2)}{4r^2(1 + r^2)^{1/2}}.$$

Therefore, the problem reduces to the family of first-order systems

$$\begin{cases} \dot r = \rho \\ \dot\rho = r(1 + r^2)\left[\frac{d_1^2}{4m^2r^4} - \frac{d_2^2}{m^2(3+2r^2)^2}\right] + \frac{r\rho^2}{1+r^2} - \frac{m(5+4r^2)}{4r^2(1+r^2)^{1/2}}. \end{cases} \tag{96}$$

Standard results of the theory of differential equations can now be applied to system (96) to prove the existence and uniqueness of analytic negative hyperbolic Eulerian rotopulsators, for admissible initial conditions. The fixed points of system (96) are obtained from equations that reduce to polynomials, so the relative equilibria form a negligible set. This remark completes the proof. □

13. QUALITATIVE BEHAVIOUR OF THE NEGATIVE ROTOPULSATORS

In this section we will prove a result that describes the qualitative behaviour of rotopulsators in $\mathbb{H}^3$. For this purpose, we first introduce a geometric topological object that plays in $\mathbb{H}^3$ the role the Clifford torus plays for us in $\mathbb{S}^3$, and briefly present some of its properties.

13.1. **Hyperbolic cylinders.** The 2-dimensional manifold defined by

$$\mathbf{C}_{r\eta} := \{(w,x,y,z) \in \mathbb{M}^{3,1} \mid r^2 - \eta^2 = -1, \ \ 0 \le \theta < 2\pi, \ \xi \in \mathbb{R}\}, \tag{97}$$

where $w = r\cos\theta, x = r\sin\theta, y = \eta\sinh\xi$, and $z = \eta\cosh\xi$, with $r, \rho \ge 0$, is called a hyperbolic cylinder, and it has constant positive curvature for r and η fixed. But $\mathbf{C}_{r\eta}$ also lies in $\mathbb{H}^3$ because the coordinates w, x, y, z satisfy the equation

$$w^2 + x^2 + y^2 - z^2 = -1.$$

When r (and, consequently, η) take take all admissible positive real values, the family of hyperbolic cylinders thus defined foliates $\mathbb{H}^3$.

We have previously shown that negative relative equilibria rotate on hyperbolic cylinders, [6], [7]. We will next prove that, at every moment in time, a negative rotopulsator passes through a different hyperbolic cylinder of any given foliation of $\mathbb{H}^3$.

13.2. **The qualitative behaviour of negative rotopulsators.** We can now state and prove now the following result.

Theorem 2. *Consider a negative elliptic, negative hyperbolic, or negative elliptic-hyperbolic rotopulsator of the curved N-body problem in $\mathbb{H}^3$. Then, for the foliation $(\mathbf{C}_{r\eta})_{r,\eta>0}$ of $\mathbb{H}^3$ given by hyperbolic cylinders, the trajectory of each body is not contained in a single hyperbolic cylinder, but intersects a continuum of hyperbolic cylinders. For negative elliptic, negative hyperbolic, and negative elliptic-hyperbolic relative equilibria, however, each body moves on some hyperbolic cylinder of the foliation for all time.*

Proof. The qualitative behaviour of negative relative equilibria in $\mathbb{H}^3$, including more specific properties, was previously described and proved to take place, [6], [7], so we will further focus on negative rotopulsators. So consider the foliation $(\mathbf{C}_{r\eta})_{r,\eta\ge 0}$ of $\mathbb{H}^3$ with hyperbolic cylinders, where $\mathbf{C}_{r\eta}$ is defined in (97). The negative elliptic-hyperbolic rotopulsators are given by (5), with r and η nonconstant functions satisfying the relation $r^2 - \rho^2 = -1$. Each hyperbolic cylinder in the foliation, however, corresponds to a constant r and a constant η that also satisfy the relation $r^2 - \eta^2 = -1$. Consequently, at every moment in time, each particle passes through a different hyperbolic cylinder, thus travelling through a continuum of such objects during its motion.

In the case of negative elliptic rotopulsators, the rotation takes place only with respect to the wx-plane, so we can treat this situation as a particular form of negative elliptic-hyperbolic rotopulsators, for which the result was proved above. Similarly, the negative hyperbolic case can be viewed as a particular for negative elliptic-hyperbolic rotopulsator, for which the rotation takes place only relative to the yz-plane. This observation completes the proof. $\square$

14. Final remarks

Criteria 1, 2, 3, 4, and 5 open the possibility to study large classes of rotopulsators for specific values of $N \geq 3$. Since the examples we provided in this paper are restricted to $N = 3$ and regular configurations, it would be interesting to know whether there exist rotopulsators given, for instance, by irregular polygons for equal or nonequal masses. Other geometric shapes could also be investigated.

We derived the equations of motion for the positive elliptic, positive elliptic-elliptic, and negative elliptic Lagrangian rotopulsators and for the negative hyperbolic and negative elliptic-hyperbolic Eulerian rotopulsators in the 3-dimensional case, but used them mainly to obtain existence and uniqueness results. It would be interesting to study these equations in detail and provide a complete classification of these motions, as it has been done in [9] for the 2-dimensional case. Since the equations for these orbits in $\mathbb{S}^3$ and $\mathbb{H}^3$ are more complicated than the ones in $\mathbb{S}^2$ and $\mathbb{H}^2$, each system would require and extensive study. The stability of these orbits, using the tools developed in [8], is another topic that merits close attention.

Finally, the nonstandard relative equilibria, presented for the first time in this paper, deserve special attention themselves. It would be interesting to identify new classes of such solutions and understand their behaviour.

Acknowledgements. Florin Diacu is indebted to Carles Simó and James Montaldi for some enlightening discussions. The authors also acknowledge the partial support of an NSERC Discovery Grant of Canada (Florin Diacu) and a University of Victoria Fellowship (Shima Kordlou).

References

[1] J. Bertrand, Théorème relatif au mouvement d'un point attiré vers un centre fixe, *C. R. Acad. Sci.* **77** (1873), 849–853.

[2] W. Bolyai and J. Bolyai, *Geometrische Untersuchungen*, Hrsg. P. Stäckel, Teubner Verlag, Leipzig-Berlin, 1913.

[3] J.F. Cariñena, M.F. Rañada, and M. Santander, Central potentials on spaces of constant curvature: The Kepler problem on the two-dimensional sphere $\mathbb{S}^2$ and the hyperbolic plane $\mathbb{H}^2$, *J. Math. Phys.* **46** (2005), 052702.

[4] F. Diacu, On the singularities of the curved N-body problem, *Trans. Amer. Math. Soc.* **363**, 4 (2011), 2249–2264.

[5] F. Diacu, Polygonal homographic orbits of the curved 3-body problem, *Trans. Amer. Math. Soc.* **364** (2012), 2783–2802.

[6] F. Diacu, Relative equilibria in the 3-dimensional curved n-body problem, *Memoirs Amer. Math. Soc.* (to appear).

[7] F. Diacu, *Relative Equilibria of the Curved N-Body Problem*, Atlantis Press, 2012.

[8] F. Diacu, R. Martínez, E. Pérez-Chavela, and C. Simó, On the stability of tetrahedral relative equilibria in the positively curved 4-body problem, arXiv:1204.5729 (32 pages).

[9] F. Diacu and E. Pérez-Chavela, Homographic solutions of the curved 3-body problem, *J. Differential Equations* **250** (2011), 340–366.

[10] F. Diacu, E. Pérez-Chavela, and J. Guadalupe Reyes Victoria, An intrinsic approach in the curved N-body problem. The negative curvature case, *J. Differential Equations* **252** (2012), 4529–4562.

[11] F. Diacu, E. Pérez-Chavela, and M. Santoprete, The N-body problem in spaces of constant curvature. Part I: Relative equilibria, *J. Nonlinear Sci.* **22**, 2 (2012), 247–266, DOI: 10.1007/s00332-011-9116-z.

[12] F. Diacu, E. Pérez-Chavela, and M. Santoprete, The N-body problem in spaces of constant curvature. Part II: Singularities, *J. Nonlinear Sci.* **22**, 2 (2012), 267–275, DOI: 10.1007/s00332-011-9117-y.

[13] W. Killing, Die Mechanik in den nichteuklidischen Raumformen, *J. Reine Angew. Math.* **98** (1885), 1–48.

[14] V. V. Kozlov and A. O. Harin, Kepler's problem in constant curvature spaces, *Celestial Mech. Dynam. Astronom* **54** (1992), 393–399.

[15] H. Liebmann, Die Kegelschnitte und die Planetenbewegung im nichteuklidischen Raum, *Berichte Königl. Sächsischen Gesell. Wiss., Math. Phys. Klasse* **54** (1902), 393–423.

[16] H. Liebmann, Über die Zentralbewegung in der nichteuklidische Geometrie, *Berichte Königl. Sächsischen Gesell. Wiss., Math. Phys. Klasse* **55** (1903), 146–153.

[17] N. I. Lobachevsky, The new foundations of geometry with full theory of parallels [in Russian], 1835-1838, In Collected Works, V. 2, GITTL, Moscow, 1949, p. 159.

[18] J.E. Marsden, *Lectures in Mechanics*, Cambridge University Press, 1992.

[19] J. Montaldi, Relative equilibria and conserved quantities in symmetric Hamiltonian systems, in *Peyresq Lectures on Nonlinear Phenomena (Peyresq 1998/1999)*, R. Kaiser and J. Montaldi, eds., pp. 239–280, World Scientific, 2000.

[20] E. Pérez-Chavela and J.G. Reyes Victoria, An intrinsic approach in the curved N-body problem. The positive curvature case, *Trans. Amer. Math. Soc.* **364**, 7 (2012), 3805–3827.

[21] E. Schering, Die Schwerkraft im Gaussischen Raume, *Nachr. Königl. Gesell. Wiss. Göttingen* 13 July, **15** (1870), 311–321.

[22] A.V. Shchepetilov, Nonintegrability of the two-body problem in constant curvature spaces, *J. Phys. A: Math. Gen.* V. 39 (2006), 5787–5806; corrected version at math.DS/0601382.

[23] A. Wintner, *The Analytical Foundations of Celestial Mechanics*, Princeton University Press, 1947.